

RECORD

Smith-Harris

667
R
200

(1958 - 1959) - 1958 - 1959
 (1958 - 1959)

1958
 1959
 1960
 1961
 1962
 1963
 1964
 1965
 1966
 1967
 1968
 1969
 1970
 1971
 1972
 1973
 1974
 1975
 1976
 1977
 1978
 1979
 1980

Collection and Field Note Book

No. 59

(Oct. 17, 1960 - March 10, 1961)

(41374 ----41455)

	<u>Pages</u>
Johnston Island.....	1-2
Specimen collected 41374.....	2
Kwajalein.....	1-7
	124-125
Specimens collected 41374--41382.....	126-141
" " 41420	2-3
" " 41420	122
Flight Kwajalein-Majuro.....	7-8
Majuro-Jaluit.....	8-9
Jaluit-Majuro.....	121-122
Jaluit.....	12-45
	114-120
Jabwor.....	12, 16
	56, 123
Specimens collected 41383--41392	18-19
Kabbenbock.....	13-15
Majurirek.....	17
Eybor.....	44-45
	56
Specimens collected 41393-41404.....	44-45
Long.....	46-55
Pinlep.....	57-65
Specimens collected 41405--41409.....	64-65

UNITED STATES DEPARTMENT OF THE INTERIOR

NATIONAL PARK SERVICE

Kinajon.....	64-65
	70-79
Specimens collected 41410--41411..	64-65
Imruj	64-65, 80-83, 97
Specimen collected 41412.....	64-65
Trip Jabwor to Imruj.....	66-69
Ligeron.....	84-86
Mejatto.....	87-96, 102-113
Specimens collected 41414--41419	
Ribon.....	98-101
Naru.....	124
Hawaiian Islands.....	142-143
Specimens collected 41421--41422	142-143
" " 41429--41442	168-171
" " 41443-41455	192-193
Oahu.....	152-200
Flight Honolulu to San Francisco....	142-143
California.....	144-145
Virginia	
Fairfax County.....	146-149
Chantilly Airport	147
Lake Barcroft	148-149
Specimens collected 41423-41428..	148-149
Great Falls.....	151
Falls Church, Seven Corners	150-151
Washington area.....	150-151

(see above)... 152-200
(Hawaii-Oahu)

You may
a you by
gh better

reference to

JULY 1, 1961

The system of 30 National Parks contains areas of highest scenic and scientific grandeur essentially in the primitive state. The National Park System, of the Department of the Interior, administers these, as well as 150 other areas of outstanding national significance. The law of the land enjoins that these be so used that they may be passed unimpaired for the enjoyment of future generations.

The story of HAWAII NATIONAL PARK is the story of active volcanism distinguished by eruptions of very fluent lava. The park is in two sections: that on the island of Hawaii embraces the summit of Mauna Loa and most of Kilauea Volcano; the section on the island of Maui includes the great crater of Haleakala.

In the maze of lava streams that pour out during Hawaiian eruptions, areas of older surface often escape cover by successive flows. An oasis thus formed is called a *kipuka*, a word that also means a loop or an open space in the forest. If a *kipuka* of large size is effectively isolated for a long period of time, good, deep soil forms, the plant inhabitants develop characteristics that reflect the nature of their history and environment, a distinctive flora is evolved. Such is the *Kipuka Puau* with its area of 100 acres. Some forty-odd species of native trees have been claimed as natural members of its flora. Many type specimens have been collected in it and described by botanists. Authorities have been drawn from faraway countries to observe its treasures. It is a significant place in the world of plants.

The name has been interpreted variously. *Pua* means flower, but is also to be translated as a collection of things bound together. *Ulu* is to grow; it is also the name for breadfruit, which appears inapplicable here. A popular nickname, Bird Park, has no significance and little to commend it.



Mamane
See No. 8



Pilo
See No. 13

To label a plant or natural specimen is too often prelude to its destruction or removal. Please respect the integrity of specimens and be thoughtful of those who follow you.

Cigarette butts and trash do not add to your enjoyment of the park. You can easily dispose of them so that they cannot be obnoxious to you and to those who come after you.

At the start of the trail is an exhibit with a map showing the outlines of the *kipuka*, surrounded by prehistoric but recent (before 1800) flows. Four common native birds, the *apapane*, *iiwi*, *ama-kihi*, and *elepaio*, are illustrated in color. This is a favorite haunt, giving the nickname *Bird Park*. Since the birds forage in the lofty canopy of trees, and sing little during the heat of the day, patience and necessary qualities for seeing and studying these alluring subjects.



Aalii with capsules

Please help preserve the primitive scene by placing refuse in containers provided for this purpose and by refraining from picking, breaking, or removing flowers, ferns, and natural specimens.

Dodonaeas are found on all of the larger islands of Hawaii. The three chief species are *D. viscosa*, *D. eriocarpa*, and *D. sandwicensis*. *D. viscosa* is also found in other parts of the Pacific.

2. Ohia Lehua (*Metrosideros collina*). The *ohia*, like the eucalyptus and guava, belongs to the Myrtle Family. It is found throughout Polynesia, and grows from sea-level to 9,000 feet in Hawaii. It is the commonest forest tree on the islands often growing in almost pure stands. It is a pioneer on most lava flows. The showy pompoms of bright red stamens make it attractive while in flower.

3. As you stand on this rise, look about you. You are on the threshold of Kipuka Puau. Note the contrasting scenes behind and before you. Almost all of the trees on the flows back of you are *ohia*; the shrubs mixed through them are *pukiawe* (*Styphelia tameiameia*) and *aalii*.

4. Ionui (*Dryopteris paleacea*). A rather stiff, erect fern that is common in Bird Park. Varieties or close relatives of this fern are widespread throughout the world. This and several other ferns are erroneously termed *laukahi*, the Hawaiian word for plantain, *Plantago major*.

5. Palapalai (*Microlepia setosa*), a lacy, attractive fern, one of the commonest along the trail. It prefers open glades and edges of the woods. It grows on all of the larger islands, and elsewhere on Pacific Islands, Malaysia, India, and Ceylon. It was used to decorate the altars of Laka, goddess of the hula.

The system of 30 National Parks contains areas of highest scenic and scientific grandeur essentially in the primitive state. The National Park System, of the Department of the Interior, administers these, as well as 150 other areas of outstanding national significance. The law of the land enjoins that these be so used that they may be passed unimpaired for the enjoyment of future generations.

The story of HAWAII NATIONAL PARK is the story of active volcanism distinguished by eruptions of very fluent lava. The park is in two sections: that on the island of Hawaii embraces the summit of Mauna Loa and most of Kilauea Volcano; the section on the island of Maui includes the great crater of Haleakala.

In the maze of lava streams that pour out during Hawaiian eruptions, areas of older surface often escape cover by successive flows. An oasis thus formed is called a *kipuka*, a word that also means a loop or an open space in the forest. If a *kipuka* of large size is effectively isolated for a long period of time, good, deep soil forms, the plant inhabitants develop characteristics that reflect the nature of their history and environment, a distinctive flora is evolved. Such is the *Kipuka Puau* with its area of 100 acres. Some forty-odd species of native trees have been claimed as natural members of its flora. Many type specimens have been collected in it and described by botanists. Authorities have been drawn from faraway countries to observe its treasures. It is a significant place in the world of plants.

The name has been interpreted variously. *Pua* means flower, but is also to be translated as a collection of things bound together. *Ulu* is to grow; it is also the name for breadfruit, which appears inapplicable here. A popular nickname, Bird Park, has no significance and little to commend it.



Mamane
See No. 8



Pilo
See No. 13

To label a plant or natural specimen is too often prelude to its destruction or removal. Please respect the integrity of specimens and be thoughtful of those who follow you.

Cigarette butts and trash do not add to your enjoyment of the park. You can easily dispose of them so that they cannot be obnoxious to you and to those who come after you.

At the start of the trail is an exhibit with a map showing the outlines of the *kipuka*, surrounded by prehistoric but recent (before 1800) flows. Four common native birds, the *apapane*, *i'iwi*, *ama-kihī*, and *elepaio*, are illustrated in color. This is a favorite haunt, giving the nickname *Bird Park*. Since the birds forage in the lofty canopy of trees, and sing little during the heat of the day, patience and quiet are necessary qualities for seeing and studying these alluring subjects.

1. *Aalii* (*Dodonaea* sp.), one of the commonest shrubs to be seen along roads and trails in the park. It grows to the size of a small tree along the Mauna Loa Truck Trail. Male and female flowers appear on separate plants. The clusters of yellow or bright red seed capsules with papery wings make the shrub showy while in fruit. These were used for leis and for extracting a brilliant red dye with boiling water. *Kapa* and other things were colored with it. The wood of the plant is tough and durable, useful for spears and various purposes.



Aalii with capsules

Please help preserve the primitive scene by placing refuse in containers provided for this purpose and by refraining from picking, breaking, or removing flowers, ferns, and natural specimens.

Dodonaeas are found on all of the larger islands of Hawaii. The three chief species are *D. viscosa*, *D. eriocarpa*, and *D. sandwicensis*. *D. viscosa* is also found in other parts of the Pacific.

2. *Ohia Lehua* (*Metrosideros collina*). The *ohia*, like the eucalyptus and guava, belongs to the Myrtle Family. It is found throughout Polynesia, and grows from sea-level to 9,000 feet in Hawaii. It is the commonest forest tree on the islands often growing in almost pure stands. It is a pioneer on most lava flows. The showy pompoms of bright red stamens make it attractive while in flower.

3. As you stand on this rise, look about you. You are on the threshold of Kipuka Puau. Note the contrasting scenes behind and before you. Almost all of the trees on the flows back of you are *ohia*; the shrubs mixed through them are *pukiawe* (*Styphelia tameiameia*) and *aalii*.

4. *Ionui* (*Dryopteris paleacea*). A rather stiff, erect fern that is common in Bird Park. Varieties or close relatives of this fern are widespread throughout the world. This and several other ferns are erroneously termed *laukahi*, the Hawaiian word for plantain, *Plantago major*.

5. *Palapalai* (*Microlepia setosa*), a lacy, attractive fern, one of the commonest along the trail. It prefers open glades and edges of the woods. It grows on all of the larger islands, and elsewhere on Pacific Islands, Malaysia, India, and Ceylon. It was used to decorate the altars of Laka, goddess of the hula.

6. **Sword Fern, Nianiau or Okupukupu** (*Nephrolepis exaltata*). Several species exist in the park. Native to tropical and subtropical climes, the sword fern is a popular house plant that exhibits many varieties. This is a pioneer plant in new lava fields and on the floor of Kilauea Crater. A very hairy variety, *N. hirsutula*, grows abundantly in steam cracks.

7. The common herbs forming a ground cover here are an introduced geranium (7A) and the native strawberry (7B), *ohelo papa* (*Fragaria chiloensis*). The latter bears fruits that are small, white, and often of good flavor. This strawberry is distributed from southern Chile to the Aleutians.

The vigorous grove contains *papala kepau*, *kopiko*, *pilo*, *olopua*, *kolea*, *mamane*, *mamaki*, *mane* and *ohia lehua* trees. These are discussed individually along the trail.

8. **Mamane** (*Sophora chrysophylla*). A common shrub or small tree found from 2,000 - 10,000 feet on all islands except Molokai, this member of the bean family bears attractive yellow flowers and compound leaves. It is related to the golden shower trees. It is eagerly eaten by livestock and feral goats, that have eradicated it in many areas. The wood is very hard and durable. Note the winged pods and shiny orange or yellow seeds, if the trees are in fruit. The *mamane* is endemic to the Hawaiian Islands, but other species of the genus, more than two dozen in number, are found distributed throughout warmer regions of the world.

9. **Kopiko** (*Straussia Hillebrandii*) is an abundant tree in Bird Park. It belongs to the important Coffee Family, which is widespread, is mostly tropical, and has 350 genera, of which a dozen are native to the Hawaiian Islands. Of these, *Straussia*, *Gouldia*, and *Bobea* are endemic. *Kopiko* in Bird Park are readily identified by their dark green, opposite leaves that have conspicuous midribs and veins. A scaly aphid often infests the lower surface of the leaves.



Kopiko

10. **Hawaiian olive, Olopua** (*Osmanthus sandwicensis*). The glossy, leathery leaves may be six inches long on very young trees. The hard, dense wood was used for handles of spears and adzes. *Osmanthus* belongs to the same family as the cultivated olive, *Olea europea*.



Olopua

11. **Soapberry, Manele** (*Sapindus saponaria*). It is remarkable that this tree appears to be identical with the evergreen soapberry of tropical America. Here the leaves become yellow in autumn and are shed in winter. The smooth gray bark on larger trees peels off in big flakes. The leaves are compound with four to six pairs of leaflets; on young plants they often grow on winged rachises, flattened stemlets. The pulpy seed coverings contain saponin, which foams in water. The hard, round, brownish-black seeds are sought by Hawaiians for bead leis.

12. **Papala kepau** (*Pisonia inermis*). The shiny, oblong leaves are dark and glossy. Like those of many other Hawaiian trees, they may be very large on young trees. The fruits, borne in a loose, open panicle 6 to 12 inches long, exude a viscid glue, used by the early Hawaiians for catching perching birds to obtain feathers for feather cloaks and ornaments. The word *kepau* means a viscous liquid like tar or molasses.

13. **Pilo** (*Coprosma rhyncocarpa*). Of the half hundred species in this genus, fifteen have been described from the Hawaiian Islands. Several species occur as trees and shrubs in the park; of these sprawling *kukaenene*, so common on barren lava fields and open forest, is best known. A member of the Coffee Family, little use appears to have been made of these small trees. The *pilo* is readily recognized by its opposite leaves with broad triangular stipules between their stems forming a loose, funnel-shaped sheath.

14. **Bracken, Kilau** (*Pteridium aquilinum*). The cosmopolitan bracken, possibly the best-known fern, grows abundantly in open grassland. The native form is found from elevations of 500 to 9,500 feet. The plant has a creeping, underground stem upon which the large, coarse, triangular fronds grow. Varied use is made of the plant throughout the world such as for food, medicine, litter, and basket material. The fruiting bodies of bracken occur in a continuous band along the edges of a frond, which are partially rolled back over it.



Papala kepau

6. **Sword Fern, Nianiau or Okupukupu** (*Nephrolepis exaltata*). Several species exist in the park. Native to tropical and subtropical climes, the sword fern is a popular house plant that exhibits many varieties. This is a pioneer plant in new lava fields and on the floor of Kilauea Crater. A very hairy variety, *N. hirsutula*, grows abundantly in steam cracks.

7. The common herbs forming a ground cover here are an introduced geranium (7A) and the native strawberry (7B), *ohelo papa* (*Fragaria chiloensis*). The latter bears fruits that are small, white, and often of good flavor. This strawberry is distributed from southern Chile to the Aleutians.

The vigorous grove contains *papala kepau*, *kopiko*, *pilo*, *olopua*, *kolea*, *mane*, *mamaki*, *mane* and *ohia lehua* trees. These are discussed individually along the trail.

8. **Mamane** (*Sophora chrysophylla*). A common shrub or small tree found from 2,000-10,000 feet on all islands except Molokai, this member of the bean family bears attractive yellow flowers and compound leaves. It is related to the golden shower trees. It is eagerly eaten by livestock and feral goats, that have eradicated it in many areas. The wood is very hard and durable. Note the winged pods and shiny orange or yellow seeds, if the trees are in fruit. The *mamane* is endemic to the Hawaiian Islands, but other species of the genus, more than two dozen in number, are found distributed throughout warmer regions of the world.

9. **Kopiko** (*Straussia Hillebrandii*) is an abundant tree in Bird Park. It belongs to the important Coffee Family, which is widespread, is mostly tropical, and has 350 genera, of which a dozen are native to the Hawaiian Islands. Of these, *Straussia*, *Gouldia*, and *Bobea* are endemic. *Kopiko* in Bird Park are readily identified by their dark green, opposite leaves that have conspicuous midribs and veins. A scaly aphid often infests the lower surface of the leaves.



Kopiko

10. **Hawaiian olive, Olopua** (*Osmanthus sandwicensis*). The glossy, leathery leaves may be six inches long on very young trees. The hard, dense wood was used for handles of spears and adzes. *Osmanthus* belongs to the same family as the cultivated olive, *Olea europea*.



Olopua

11. **Soapberry, Manele** (*Sapindus saponaria*). It is remarkable that this native tree appears to be identical with the evergreen soapberry of tropical America. Here the leaves become yellowish in autumn and are shed in winter. The smooth gray bark on larger trees peels off in big flakes. The leaves are compound with four to six pairs of leaflets; on young plants they often grow on winged rachises, i.e., flattened stemlets. The pulpy seed coverings contain saponin, which lathers in water. The hard, round, brownish-black seeds are sought by Hawaiians for bead leis.

12. **Papala kepau** (*Pisonia inermis*). The shiny, oblong leaves are dark and glossy. Like those of many other Hawaiian trees, they may be very large on young trees. The fruits, borne in a loose, open panicle 6 to 12 inches long, exude a viscid glue, used by the early Hawaiians for catching perching birds to obtain feathers for feather cloaks and ornaments. The word *kepau* means a viscous liquid like tar or molasses.



Papala kepau


13. **Pilo** (*Coprosma rhyncocarpa*). Of the half hundred species in this genus, fifteen have been described from the Hawaiian Islands. Several species occur as trees and shrubs in the park; of these sprawling *kukaenene*, so common on barren lava fields and open forest, is best known. A member of the Coffee Family, little use appears to have been made of these small trees. The *pilo* is readily recognized by its opposite leaves with broad triangular stipules between their stems forming a loose, funnel-shaped sheath.

14. **Bracken, Kilau** (*Pteridium aquilinum*). The cosmopolitan bracken, possibly the best-known fern, grows abundantly in open grassland. The native form is found from elevations of 500 to 9,500 feet. The plant has a creeping, underground stem upon which the large, coarse, triangular fronds grow. Varied use is made of the plant throughout the world such as for food, medicine, litter, and basket material. The fruiting bodies of bracken occur in a continuous band along the edges of a frond, which are partially rolled back over it.


KILAUEA CRATER

VICINITY MAP

Roads, Trails, and Points of Interest



The Park Museum is in the Park Administration Building on the main highway, one mile inside the Hilo Entrance to the Park. It is open from 7:30 a.m. to 4:30 p.m. daily, and is dedicated to the wonders of the park story. Motion pictures of eruptions of Mauna Loa and Kilauea are shown daily in the Park Auditorium in the same building at 9:30 a.m. and 1:15 and 1:45 p.m. Admission is free. Information is dispensed at the counter in the lobby of the Administration Building; maps, copies of motion pictures and kodaslides, and literature are on sale at this place by the Association. All profits are used for furthering interpretive efforts of the park.



KEY

- Cinder and spatter cones and ramparts at vents
- Trace of eruptive fissure
- Zones of cracking formed during the 1954 eruption

0 5000 feet



UNITED STATES
DEPARTMENT OF INTERIOR
Fred A. Seaton, Secretary
NATIONAL PARK SERVICE
Conrad L. Wirth, Director

15. **Manena** (*Pelea cinerea*) has slender, twining branchlets with small leaves and comparatively small fruits. Like the others of its genus, the leaves are opposite; some are fragrant.

16. **Mamaki** (*Pipturus sp.*), is a member of the stinging nettle family. The inner bark of this plant was used by the Hawaiians for making *kapa* or bark cloth. It is one of the most abundant fiber plants found in the Islands.

17. **Manono** (*Gouldia terminalis*). The lone tree ten paces above the sign is another member of the Coffee Family. The genus is one of the three endemic to the Hawaiian Islands.



Manono

18. **Kolea** (*Myrsine sp.*). The red color of the young leaves is an interesting feature of this plant. It is due to the chemical anthocyanin that protects new, tender leaflets from the brilliant sun of the tropics. Wood from this tree was used for houseposts; a dye for *kapa* was made from the sap. *Kolea* is also the name given to the golden plover, the bird that completes a remarkable migration to the Arctic Circle and back to Hawaii each year.

19. **Nasturtium** (*Tropaeolum majus*), a native of Peru, that has escaped from gardens, finding the environment very suitable for its spread. Here it drapes a prostrate *ohia* log, ten paces below the trail.

20. **Koa** (*Acacia koa*). The trees in the glade below the trail are *koa*. Like the *mamane*, the *koa* is a member of the legume or pea family and is native only to the Hawaiian Islands. The genus *Acacia* has 450 species in the tropics and subtropics throughout the world; those growing in Australia, are called wattles. The bark of the young *koa* is smooth; the bark on the older trees is coarse.

21. **Maua** (*Xylosma Hillebrandii*). This small tree is ten paces behind the sign. It is uncommon in Bird Park, but prefers drier lands on leeward slopes of all the major islands except Kauai and Oahu. It is a handsome tree with shiny, toothed, crenulate leaves whose color often gives the tree a reddish cast.

22. **Kipuka Boundary**. This ridge is the edge of the prehistoric lava flow from Mauna Loa that surrounded this *kipuka* but failed to cover it. Notice the difference between the forest growing on the relatively recent lava flow ahead and that growing on the rich soil of the *kipuka*.



Maile No. 36

23. **Papala** (*Charpentiera obovata*), an endemic tree, rare in the park. Do not touch or disturb this planted specimen. Its small red flowers hang in large loose panicles from the ends of branches. The wood is soft, fibrous, very light, and inflammable. The Hawaiians heaped it into bonfires which they pushed over cliffs and vantage points. Because of their buoyancy, rising air currents would buffet the glowing torches in brilliant display upward, downward, and sideways like so many shooting stars.



Papala

24. **Alaalawainui** (*Peperomia reflexa*). A large number of species of this widespread genus of the Pepper Family grow in Hawaii. Condiment pepper is made from *Piper nigrum*, native to the Oriental tropics. The celebrated Polynesian drink, *awa* or *kava*, was made from the root of *Piper methysticum* which grows 6 to 12 feet tall.

25. **Alani** (*Pelea volcanica*). The great botanist, Asa Gray, named this genus of the Rue (Citrus) Family after the goddess Pele. The leaves are opposite, often notched at the apex. When crushed they emit a fragrant odor. The fruiting capsules are large, usually over one inch in diameter, and open into four parts, each bearing two shiny black seeds.



Naio

26. **False Sandalwood, Naio** (*Myoporum sandwicense*). When the supply of true sandalwood (*Santalum sp.*), a chief item of trade in the early days, began to run low, *naio* was used as an inferior substitute. Since demand for it never became great, *naio* trees are still abundant in these islands, to which they are native. In Bird Park old trees develop deeply furrowed, knobby boles, with heartwood much decayed so that only strips of living tissue remain. (See illustration on page 12). Despite this, they cling tenaciously to life, sending out vigorous shoots from their tops. These are the most picturesque objects along the trail.

27. **Tarweed, Pukamole** (*Lythrum maritimum*). A sticky, hairy undershrub of the American tropics grows rank here. It bears small, magenta flowers. It belongs to the Grape Myrtle Family.



Alani



Geranium

28. Open Forest Growth. In this locale, the trees do not compete with each other in their efforts to obtain sunlight. Compare the symmetry of the spreading ohia in the distance to the crowded *koa* on the left skyline. Note also the fallen *ohia* twenty paces to the left. Its branches have turned upward and in time may become independent trees as the old tree decays. Curiously shaped *ohia* may be seen on the right of the trail beyond sign 39, and on the left of the trail 30 paces before reaching sign 42. Similar contorted individuals are especially conspicuous at Kipuka Nene campground.

This is an excellent spot for observing birds. The frequent crowing of pheasant cocks is a complementary sound.

29. Collapsed Lava Tube. Many layers of volcanic ash have accumulated through the centuries, some of them from eruptions of distant Mauna Kea. Such ash forms much of the topsoil in Kipuka Puau. This collapsed lava tube and the lava above it provide opportunity for gauging the depth of the soil. **DANGER—DO NOT APPROACH THE TUBE.**

30. Kookoolau (*Bidens pilosa*), an introduced weed from tropical America. Its annoying, black needles, less than one-half inch long, are pronged so that they readily cling to clothes and to fur of animals. The many species of *Bidens* have been given appropriate popular names: Spanish needles, beggar ticks, stickights, tick seed, bur marigolds. Over sixty blander native Hawaiian species have been described. Leaves and tips of young plants, fresh or dried, are steeped for a beverage, often in preference to commercial tea. In season, a large undershrub native to Puuwaawaa glitters with thousands of golden flowers in big panicles, a glorious sight indeed. Kookoolau Crater on Chain of Craters Road was named for the species growing within it.

31. Ae (*Zanthoxylum dipetalum*). The *Zanthoxyla*, represented by three species in Bird Park, are very rare in numbers of individuals. This is an unusually large specimen. Like the *Pelea*, this belongs to the Rue (Citrus) Family.



Ae leaf

32. Akala (*Rubus hawaiiensis*), a giant among raspberries. A half dozen berries fill a bowl. The bearing season, which is in midyear, like the taste and color of the fruit, varies with plants and location. Usually the berries are purple, but the blossoms and juice are pink. *Akala* means *pink* in Hawaiian.

A short trail to the left leads to a huge *koa* whose base is seven feet in diameter.



Olapa

33. Huehue (*Cocculus ferrandianus*), a native climber widely distributed from sea coast to 5,000 feet. The tough flexible stems are useful as natural twine. Hawaiians wove them into baskets with funnellform mouths that were used as fish traps.

34. Alani (*Pelea Zahlbruckneri*). Named by the great Hawaiian dendrologist, Joseph Rock, for his celebrated friend, Dr. A. Zahlbruckner, Director of the Viennese Botanical Museum. It has the largest leaves of any *Pelea*, and large, curious fruiting capsules like four-pointed stars.

35. Olapa (*Cheirodendron Guadichaudii*). One of the commonest, most conspicuous of forest trees. Its bright green, shiny leaves are palmately compound and constantly flutter in the breeze. When bruised, all parts of the tree emit a strong turpentine odor. The name *olapa* was applied to those hula dancers with lithe, supple bodies and most graceful motions, who could best imitate the dancing of the *olapa* leaves. The tree is twenty paces behind this sign, growing in the big patch of *Microlepia*.

36. Maile (*Alyxia olivaeformis*). This twining, native, vinelike shrub is the laurel of old Hawaii. Its fragrant stems and shiny leaves carry a faint odor of vanilla. They were a favorite and an indispensable adjunct to every festive occasion, being used for decoration and for leis.

37. Mint (*Mentha spicata*). This large patch of non-native mint affords an example of how native plants are often crowded out by hardy foreigners. Please do not pick the mint and drop it along the trail, as this aids in spreading it and in choking out desirable plants.

38. Large Koa (*Acacia koa*). In regions in which growing seasons are indefinite, the age of a tree cannot readily be determined by a count of annular rings. This tree is probably several hundred years old. The *koa* was used more than any other tree in making Hawaiian canoes, both the single *kaukahi* and the double *kaulua*. The hard, beautiful wood is suitable for the manufacture of furniture and other objects, such as bowls and trays.

39. Opuhe (*Urera sandwicensis*), like the *mamaki*, is a member of the stinging nettle family and is used for making *kapa*. It is a medium-sized tree with male and female flowers growing on separate plants. The large, oblong, dark green leaves have prominent veins; the stems exude a watery, milky fluid when broken. Fibers from the bark are tough, useful for making cord for fish nets.

40. Large Ohia Lehua (*Metrosideros collina*). An idea of the size to which these trees grow may be obtained from this specimen. It is approximately 80 feet high, and about five feet in diameter at the base. The flower of the *ohia*, the *lehua*, is the flower of the Island of Hawaii. Many birds, particularly the red *apapane*, feed on the nectar of these flowers. The wood of *ohia* is hard and durable; it was used for making gunwales of outrigger canoes, timbers for housing, and poi boards.

41. **Ti, Ki** (*Cordyline terminalis*), is a very useful plant to Hawaiians. The leaves are used in wrapping food, either for storage, for handling, or for cooking in the earth. They formerly served for thatch and for skirt material. A crude musical instrument is made by rolling up a leaf. *Okolehao*, a potent alcoholic beverage, is brewed from the roots of this plant.




Naio trunk

42. **Young Koa Leaves** (*Acacia koa*). Both the fernlike, juvenile leaves and the adult, sickle-shaped phyllodea may be seen on the lower branches of these trees. A young *koa* starts growth with compound leaves only. As it grows, the petioles (leaf-stalks) expand and replace the leaflets. The resulting leaflike organ, termed a *phyllode*, may be a clue to the evolutionary history of the *koa*. Finely divided leaves are characteristic of plants that are not exposed to loss of life-sustaining moisture through surface evaporation. A leathery, smaller-surfaced appendage is better adapted to withstand the strain of drought. Despite heavy rainfall in the islands, supplies of moisture stop abruptly with a shift of winds: the soil and atmosphere quickly assume a desert character which, if of long duration, is critical to poorly adjusted plants. The *koa* has become neatly adjusted to its environment.

43. **Grassland.** Many grasses not native to the island grow in Kipuka Puaulu. Most of these were introduced with stock feed brought from the mainland. In some areas the grasses are growing so densely that it is impossible for native plants to get a start.

44. **Avocado** (*Persea americana*). Somebody thoughtlessly tossed aside a seed. Well above optimum range for the plant, it has grown into a robust tree that cannot bear fruit.




Alaalawainui

DESERT MUSEUM

Tucson Area

AT THE DESERT MUSEUM
one admission charge
covers all exhibits:
Adults — 75c
Children 6 to 12 — 25c
Under Six — Free



TO
BENSON
TOMBSTONE
FORT HUACHUCA
CORONADO NAT'L. MEM.
BISBEE (OPEN PIT MINE)
CHIRICAHUA NAT'L. MON.
AND POINTS EAST

ADAPTED BY
DAVID F. CARSWELL
FROM A MAP BY
THE CITY-COUNTY
PLANNING DEPT.

Souvenir Aerial Map ARIZONA-SONORA DESERT MUSEUM

A LIVING MUSEUM

P. O. BOX 5602 TUCSON, ARIZONA

We hope the use of this aerial map will add to the pleasure of your visit to the Desert Museum. The reverse side offers a directional map designed to aid those wishing to visit other places of interest in the Tucson area.

At this museum you will see interpretive displays of living animals and plants native to the states of Arizona, U.S.A., and Sonora, Mexico, with emphasis on the Sonoran Desert contained in both states.

The Desert Museum is a non-profit educational institution supported by admissions, memberships and donations. Contributions are tax deductible.

MAKE THIS YOUR MUSEUM! Membership information is available at the desk.

(Blanton & Cole aerial photo by Don Cassidy; drafting by Red Ross)



Jan 17, 1962, U.S.A.

THE DESERT MUSEUM IS open every day of the year — from 10 A.M. to 5 P.M. (until sundown on Sundays and holidays).

ARIZONA-SONORA DESERT MUSEUM

Guide Map of Tucson Area

AT THE DESERT MUSEUM one admission charge covers all exhibits:
 Adults — 75c
 Children 6 to 12 — 25c
 Under Six — Free



TO
 PAPAGO INDIAN RESERVATION
 AJO AND OPEN PIT COPPER MINE
 ORGAN PIPE CACTUS NAT'L. MONUMENT
 PUNTO PEÑASCO (ROCKY POINT), MEX.

TO
 TUBAC PRESIDIO HIST. MON.
 TUMACACORI NAT'L. MONUMENT
 NOGALES AND OLD MEXICO

TO
 BENSON
 TOMBSTONE
 FORT HUACHUCA
 CORONADO NAT'L. MEM.
 BISBEE (OPEN PIT MINE)
 CHIRICAHUA NAT'L. MON.
 AND POINTS EAST

- KEY**
- ★ ARIZONA-SONORA DESERT MUSEUM
 - ② "OLD TUCSON" MOVIE SET
 - ③ SAN XAVIER MISSION
 - ④ MUNICIPAL AIRPORT
 - ⑤ UNIVERSITY OF ARIZONA
ARIZONA STATE MUSEUM
ARIZONA PIONEERS' HIST. SOC.
 - ⑥ RANDOLPH PARK
 - ⑦ SABINO CANYON
 - ⑧ SAGUARO NAT'L. MONUMENT
 - ⑨ COLOSSAL CAVE



DIRECTIONS
 (By GEORGE!)



(ADAPTED BY
 DAVID F. CARSWELL
 FROM A MAP BY
 THE CITY-COUNTY
 PLANNING DEPT.)

DO NOT WRITE IN THESE SPACES. DC. USA.

THE DESERT MUSEUM IS
open every day of the year —
from 10 A.M. to 5 P.M.
(until sundown on Sundays
and holidays).

ARIZONA-SONORA Guide Map of



TO
PAPAGO INDIAN RESERVATION
AJO AND OPEN PIT COPPER MINE
ORGAN PIPE CACTUS NAT'L. MONUMENT
PUNTO PEÑASCO (ROCKY POINT), MEX.

TO
TUBAC PRESIDIO HIST. MON.
TUMACACORI NAT'L. MONUME
NOGALES AND OLD MEXICO

- KEY**
- ★ ARIZONA-SONORA DESERT MUSEUM
 - ② "OLD TUCSON" MOVIE SET
 - ③ SAN XAVIER MISSION
 - ④ MUNICIPAL AIRPORT
 - ⑤ UNIVERSITY OF ARIZONA
ARIZONA STATE MUSEUM
ARIZONA PIONEERS' HIST. SOC.
 - ⑥ RANDOLPH PARK
 - ⑦ SABINO CANYON
 - ⑧ SAGUARO NAT'L. MONUMENT
 - ⑨ COLOSSAL CAVE



DIRECTIO
(By GEORGE)

change to stake out our claims on Rapa.

(I keep finding odd monocot pollen grains and must get more comparative material as soon as possible. I need Zingiberaceae (one may be a *Gurouma*), Taccaceae (I have two, but need more), Comelinaceae and Musaceae.... *Maritima* isn't too good at sending samples at present, but Eddie Bryan would help, he says. He might pick out a few scurps and send them along with you. All he has sent so far has been most useful. Annatta Carter (Berkeley) has also sent pollen of 13 spp. Unfortunately most of her Rapa material--which seems to be extensive--has no pollen.

Can you get any samples of flowers of *Balanophora*? This may be important. *Ary Balanophora*--the more the better.. I have *B. fungosa*, but must check that as I believe Croizat lists a different species for Rapa.

The Rubiaceae may be important: I think that family should be treated

Jan 25, DC. USA.

Mrs. WATSON SMITH
5045 EAST GRANT ROAD
TUCSON, ARIZONA

Sunday

I have a sketch of N.E. and Hawaiian bog formers (Oracholus and Donatia) if you wished to use it... It was prepared for Verdoorn when he planned to bring out a book on Fest Deposited the world. It has never been printed. L.

Dear Ray:

Al Mead rang a short time ago to say that you would come at Easter time. He wasn't too sure whether you would arrive on A. He said he would write you and arrange to meet you at the airport. I told him to bring you here. You can have the guest-room, if you don't mind one in a Museum. It has all be cleaned up for friends due today, and others due from Malaya at the end of April. I cannot offer much except talk, but I think that is what you have come for. We can see that you get in to see Al or Paul Martin and so make good use of your time.

You have not heard from me because I have been sick and am in the frustrated frame of mind that makes me wish to underline everything. I am wholly better this week, but have had recurrent influenza all of Jan. and Feb. Never had anything like it: next Feb. I think we'll go away to a more equable climate and dodge this odd spring climate here. We had snow on the mts. today, after 84° temperatures two days ago: that is typical. I think I got run down before Xmas, and tried to do too much with the microscope at night.

I got a friend to make prints of the Hawaiian negatives, but they are only so-so. I enclose some. I could still send you the negatives, of course. Unfortunately my helper lost my treasured negative of Selling stuck in a mucky path on Kauai: that has been bad luck.

Congress: I sent in the proposed arrangement, changing the title to:--
POLLEN CLUES TO ANCIENT PACIFIC FLORAS. That has seemed to suit everyone better. I found that nobody wished to speak on Cretaceous angiosperms. I was premature about that, but plan to stick to them myself. Dr. A.C. Smith says he has sent off the title to Honolulu, so you may know all this... Jane Gray is to do Pacific NW coast; a Jap

to Honolulu, so you may know all this... Jane Gray is to do Pacific NW coast; a Jap
that, but plan to stick to them myself. Dr. A.C. Smith says he has sent off the title
found that nobody wished to speak on Cretaceous angiosperms. I was premature about
POLLEN CLUES TO ANCIENT PACIFIC FLORAS. That has seemed to suit everyone better. I
Congress: I sent in the proposed arrangement, changing the title to:--
path on Kauai: that has been bad luck.

Fortunately my helper lost my treasured negative of Selling stick in a mucky
only so-so. I enclose some. I could still send you the negatives, of course. Un-
I got a friend to make prints of the Hawaiian negatives, but they are

do too much with the microscope at night.
two days ago: that is typical. I think I got run down before Xmas, and tried to
this odd spring climate here. We had snow on the mts. today, after 8th temperatures
like it: next Feb. I think we'll go away to a more equable climate and dodge
week, but have had recurrent influenza all of 'em and Feb. Never had anything
frame of mind that makes me wish to underline everything. I am wholly better this
You have not heard from me because I have been sick and am in the frustrated

Martin and so make good use of your time.
that is what you have come for. We can see that you get in to see Al or Paul
from Malaya at the end of April. I cannot offer much except talk, but I think
one in a Museum. It has all be cleaned up for friends due today, and others due
told him to bring you here. You can have the guest-room, if you don't mind
He said he would write you and arrange to meet you at the airport. I have
at Easter time. He wasn't too sure whether you would arrive on April 2 or 3.
Al had rang a short time ago to say that you would come through

Dear Ray:



Sunday

Mrs.
WATSON SMITH
5045 EAST GRANT ROAD
TUCSON, ARIZONA

something about genera that survive in Japan(it turned out he could not tackle fossil material), Muller will contribute a paper--to be read by Jane Gray--on pollen evidence to do with the history of the mangrove formation -- and then Balme of W.Australia(if he can come) will give a survey of pollen and spore work for older deposits of Australia. I end up with "THE CRETACEOUS ROLE OF NOTHOFAGUS".

It will thus be a sort of progress report on pollen work in the Pacific.

To my chagrin Dr. Smith wrote last week to say that he thought I wanted a four hour session. I have been told, I understood, to arrange one of 2 hours with 3-4 speakers. He left the matter open, so I wrote back at once accepting the longer time and explaining that it would help greatly in developing the arguments. Chaney will now have more time to introduce to lead comments after papers or at the end, and I have suggested that he give his introductory comments a title* something like a "Macropalaeontologist looks at Fossil Pollen". He can be as critical or as appreciate as he likes and thus inform the audience about what is going on in general in palaeontology. He has been most co-operative so far, and all will be well, if he turns up. I have heard that he can change plans without much warning.

He wants to add in a Dr. Tokunaga from Japan. He has offered to head his paper--on Eocene pollen from Japan, so I have now accepted. He says neither Tokunaga nor Ueno(the other man, whom he recommended first and whom, unfortunately, we invited) should try to deliver papers in English).

Thus there is some chance to compare the northern and southern deposits and microfossils over a considerable period....The mangroves will be of general interest as a lot of attention is now being paid to them in Europe, in this country, and by Croizat. The southern hemisphere is a little short just now, and I am worried as Balme has not written since term began at the beginning of March. He had had no word then about getting any financial aid. Without aid from the N.S.F. he is in a bad position in seeking aid from his University or his Govt...He went to India two years ago and is afraid the Univ. won't help now, as they are so short of funds. He suggests very aimably that I could read his paper if he failed to turn up, but it is rather beyond me. He is a to-notcher, and most highly spoken of by Dr. Edna Plumstead, who is just about to come out with a striking contribution on the older fossil deposits from which the Fuchs-Hillary Exped. collected....You

should watch for her papers on Glossopteris too: they are used by Delville in his theories about the devt. of the angiosperm flower (Nature, Oct. 1, 1961: long article). (Or was it Feb. 1: I cannot remember). She agrees with me about Antarctica but has far more evidence than I have had for somewhat revolutionary views. I have offered her some of my new evidence, and have told her I intend to send a note in to Nature about it as soon as possible.

With two hours extra in hand, we may be in difficulties, if Balme does not come, so I have reinvited Dr. Isabel Cookson by cable, and have also asked Vishn-Mittre (Birbal Sahni Institute) by cable. I am waiting for their replies now. Cookson might come, as she indicated too late that she would have accepted my first invitation if there had been any suggestion that help would be available. Vishnu-Mittre might be able to come, as he has been asking questions about it for some time. Florin backs him very strongly, so I have asked him to do a short paper on trisaccate grains: this brings in the Pacific area, Antarctica, and Kerguelen; and then the Gondwana areas. He has plenty to say as he has been telling me his views in recent letters/

If neither can come I am afraid I could not have more papers read, as that would throw too much on Chaney, Jane Gray and myself.

This is a suggestion--not just an alternative: Would you be willing to collaborate with Stella Peopold and myself in giving a piece on Rapa? I did not invite Stella originally as I did not know of her work on the western borings, nor that she had that Rapa sample. I think I shall telephone her and get an opinion from her. I'd then ask her to deliver the paper, if she could come to the Congress. She may, of course, be going to the INQUA one in Poland.....If she cannot do it, perhaps you and I could do it, and you could deliver it. The pollen side would be slight, comparatively, and you could handle it if I got slides ready. By then you will have the note in Nature to refer to. (I am sure they will accept it, as they like controversial things). PLEASE SEND ME A CABLE OR AIRMAIL LETTER AS SOON AS POSSIBLE, AS SMITH GIVES ME ONLY UNTIL MARCH ~~1st~~ 31st. to inform him about full list of speakers, titles, times, summaries, etc. I think this would be a

chance to stake out our claims on Rapa.

(I keep finding odd monocot pollen grains and must get more comparative material^a as soon as possible. I need Zingiberaceae (one may be a Curcuma), Taccaceae (I have two, but need more), Commelinaceae and Musaceae.....Marie Neal isn't too good at sending samples at present, but Eddie Bryan would help, he says. He might pick out a few scarps and send them along with you. All he has sent so far has been most useful. Annetta Carter (Berkeley) has also sent pollen of 13 spp. Unfortunately most of her Rapa material--which seems to be extensive-- has no pollen.

Can you get any samples of flowers of Balanophora? This may be important. Any Balanophora--the more the better.. I have B. fungosa, but must check that as I believe Croizat lists a different species for Rapa.

The Rubiaceae may be important: I think that family should be treated monographically (for pollen types)...

The total of species for Rapa is still not high, so don't let me raise your hopes too much. This is odd, somehow, as some delicate grains are marvellously well-preserved. Bailey did not agree that my tracheids were from an Araucarian: I have lost faith in identifying this family from wood elements, anyway: most of the literature is conflicting....

The conifer pollen remains, but it is now more difficult to handle. You will see what I have done about it in the Nature article: I did not attempt to hang the whole story on it, as I had hoped to do. The title now is (if approved by Nature) :- "RAPA NEGLECTA: A Thimbleful of Continentality". The previous one was "A Rapa Conifer: Contaminant, endemic, or Gift of the West Wind Drift?" We may still use that title/Tell me at once if you ~~prefer~~ prefer the second title.

My guests are overdue, so this letter has grown longer.

Best wishes,

Lacey

P.S. If you send a cable you can use our Tucson Tel. number: East 6.9607. That helps get it to use quickly. Address it to me as MRS. WATSON SMITH.

LES

monographically (for pollen types)...

The total of species for Rapa is still not high, so don't let me raise your hopes too much. This is odd, somehow, as some delicate grains are remarkably well-preserved. Bailey did not agree that my tracheids were from an Araucarian; I have lost faith in identifying this family from wood elements, anyway; most of the literature is conflicting....

The conifer pollen remains, but is now more difficult to handle. You will see what I have done about it in the Nature article; I did not attempt to hang the whole story on it, as I had hoped to do. The title now is (if approved by Nature) "RAPA NEGIOTIA: A Thimbleful of Continentality". The previous one was "A Rapa Conifer"; Contentment, endemic, or Gift of the "West Wind Drift?" We may still use that title/Tell me at once if you ~~approve~~ prefer the second title.

My guests are overdue, so this letter has grown longer.


Best wishes,

Harry

P.S. If you send a cable you can use our Tucson Tel. number: East 6,9507. That helps get it to me quickly. Address it to me as MRS. WATSON SMITH.

W.S.

JALUIT



4255 Survey

74

2

ing

FRANK AUBREY HICKEY
8966 DIAMOND HEAD ROAD
HONOLULU 15, HAWAII

no return
address

act

mail

DC USA.

monographically for pollen types)...

The total of species for Hapa is still not high, so don't let me
raise your hopes too much. This is not a
+ +

1.



8th NATIONAL
WATERSHED CONGRESS

sales
all in

374



- ① For Road
with 11 acres
Substation
- ② to be
acquired by
Mansp. comm.

set
mail

7th and 15th, DC. USA.

8
III NATIONAL
WATERSHED CONGRESS

RAMADA INN

TUCSON, ARIZONA

APRIL 17, 18, AND 19, 1961



THEME: COUNT DOWN ON WATER

PROGRAM

It is the intention of the National Park Service to further scientific research within the areas administered by it, and to cooperate with technical workers to the fullest extent compatible with its charge to preserve all species of flora and fauna and all geologic material in a natural state, insofar as is possible.

CONDITIONS UPON WHICH THIS COLLECTING PERMIT IS ISSUED

74

e
ing

object
unmail
USA.

THEME: COUNT-DOWN ON WATER

PROGRAM

SUNDAY, APRIL 16

- 3:00 p.m. Registration
Lobby, Ramada Inn
- 8:00 p.m. Special Motion Picture Program
Ramada Room

MONDAY, APRIL 17

- 8:00 a.m. Registration
Lobby, Ramada Inn

**SESSION 1
Ramada Room**

Theme: "Planning for Water Use"

- 9:50 a.m. Formal Opening
C. R. Gutermuth, Chairman, Steering Committee, National Watershed Congress
 - 10:00 a.m. Water Needs of the Nation 1980-2000
Hon. Robert S. Kerr
United States Senator from Oklahoma
 - 10:30 a.m. Discussion
 - 10:40 a.m. Soil and Water Conservation Needs Inventory
Donald A. Williams, Administrator, U. S. Soil Conservation Service, Washington, D. C.
 - 11:10 a.m. Discussion
 - 11:20 a.m. Legal Barriers in Water Use Planning
Frank J. Trelease, Dean, College of Law, University of Wyoming, Laramie
 - 11:50 a.m. Discussion
 - 12:00 noon Recess
 - 12:15 p.m. Luncheon, **Plantation Room**
James B. Craig, Presiding
- Address of Welcome
Arthur N. Pack, President, Charles Lathrop Pack, Forestry Foundation, Tucson
- Development of "Water Street U.S.A."
William H. Carr, Founder and Director Emeritus, Arizona-Sonora Desert Museum, Tucson

SESSION 2

- 2:00 p.m. Buses leave Ramada Inn for meeting on "Water Street U.S.A."
- 5:30 p.m. Buses leave for "Old Tucson"
- 6:00 p.m. Reception, chuck wagon dinner, and entertainment at "Old Tucson"

TUESDAY, APRIL 18

- 8:00 a.m. Registration
Lobby, Ramada Inn

**SESSION 3
Ramada Room**

Theme: "Multiple Use of Watersheds"

- 9:00 a.m. William E. Richards, Presiding
 - 9:00 a.m. Opportunities in Multiple Use of Watersheds
Harold G. Wilm, Commissioner, New York State Conservation Department, Albany
 - 9:30 a.m. Discussion
 - 9:50 a.m. Water for Agriculture, Industry and Recreation
Hon. Frank J. Welch, Assistant Secretary of Agriculture, Washington, D. C.
 - 10:20 a.m. Discussion
 - 10:40 a.m. Motivating Multiple Use of Watersheds
Marvin Melton, President, Arkansas State Chamber of Commerce, Little Rock
 - 11:10 a.m. Discussion
 - 11:30 a.m. Recess
 - 11:45 a.m. Luncheon, **Plantation Room**
John H. Jones, Presiding
- Life Zones in Evidence on Spectacular Mt. Lemmon
Norman P. Weeden, Supervisor, Coronado National Forest, Tucson
- 1:30 p.m. Buses leave Ramada Inn for Mt. Lemmon tour
 - 5:00 p.m. Buses return to Ramada Inn
 - 7:00 p.m. Reception, **Plantation Room**
 - 7:30 p.m. Annual Dinner, **Ramada Room**

Toastmaster:

C. R. Gutermuth, Chairman, Steering Committee, and Vice-President, Wildlife Management Institute, Washington, D. C.

Presentation of Awards:

Watershed of the Year
Watershed Man of the Year

Principal Speaker:

"How Small Watershed Programs Help Communities Grow"
L. L. Males, President, Security State Bank, Cheyenne, Oklahoma
"Desert Ark" live animal demonstration by Hal Gras, Tucson. Music will conclude the program

WEDNESDAY, APRIL 19

**SESSION 4
Ramada Room**

Theme: "Water for a Thirsty Land"

- John I. Taylor, Presiding
 - 9:00 a.m. Arizona Water Management Programs
Andrew L. McComb, Head, Department of Watershed Management, University of Arizona, Tucson
 - 9:30 a.m. Discussion
 - 9:45 a.m. Water Yields from National Forests
Richard E. McArdle, Chief, U. S. Forest Service, Washington, D. C.
 - 10:15 a.m. Discussion
 - 10:30 a.m. Watershed Management on Public Lands
Karl S. Landstrom, Director, Bureau of Land Management, Washington, D. C.
 - 11:00 a.m. Discussion
 - 11:15 a.m. Watershed Management to Serve Reclamation Needs
LaSelle E. Coles, President, National Reclamation Association, Prineville, Oregon
 - 11:45 a.m. Discussion
 - 12:00 noon Recess
 - 12:30 p.m. Luncheon, **Plantation Room**
William E. Richards, Presiding
- Future Water Needs for Recreation
Francis W. Sargent, Executive Director, Outdoor Recreation Resources Review Commission, Washington, D. C.
- 3:30 p.m. Special meeting for representatives of participating organizations. **Ramada Room**

THURSDAY, APRIL 20

Post-Congress tour of the Agricultural Research Service's Walnut Gulch Watershed. The 65-square-mile experimental project is located near Tombstone. This guided tour will take the greater part of the day. A luncheon stop will be made in Tombstone.

- 8:00 a.m. Buses leave Ramada Inn

HOTELS & RATES

RAMADA INN Single: \$9.00-10.00 Double: \$9.00-12.00 Twin Beds: \$9.00-12.00	TUCSON DESERT INN Single: \$8.00-10.00 Double: \$8.00-12.00 Twin Beds: \$9.00-12.00
SANDS MOTOR HOTEL Single: \$9.00-10.00 Double: \$9.00-12.00 Twin Beds: \$9.00-12.00	HOLIDAY INN Single: \$8.00 Double: \$9.00 Twin Beds: \$11.00

TO: WATERSHED CONGRESS HOUSING BUREAU

c/o Tucson Convention Bureau
P. O. Box 991
Tucson, Arizona

I plan to attend the Eighth National Watershed Congress on April 17-19, 1961.

Name Title

Organization Address

1st Choice Hotel 2nd Choice Hotel

Please reserve me a single room , Double , Twin beds for arrival on and departure on (Reservations will not be held after 6:00 p.m.)

Send confirmation to

Enclosed is \$..... for **ADVANCE REGISTRATION** for persons (at \$5 each) named below:

(Make checks payable to the Eighth National Watershed Congress)

CONGRESS STEERING COMMITTEE

C. R. GUTERMUTH
Chairman

Vice-President, Wildlife Management Institute
Washington, D. C.

JAMES B. CRAIG
Editor

American Forests
The American Forestry Association
Washington, D. C.

JOHN H. JONES
Secretary-Treasurer
American Watershed Council
Fairmont, West Virginia

WILLIAM E. RICHARDS
President
National Association of Soil Conservation Districts
Holdrege, Nebraska

JOHN I. TAYLOR
Assistant Legislative Director
American Farm Bureau Federation
Washington, D. C.

PARTICIPATING ORGANIZATIONS

- American Farm Bureau Federation
- American Fisheries Society
- American Forestry Association
- American Nature Association
- American Planning and Civic Association
- American Pulpwood Association
- American Watershed Council
- Chamber of Commerce of the United States
- Intern'l Ass'n of Game, Fish and Conservation Comm'rs
- Izaak Walton League of America
- National Association of County Officials
- National Association of Manufacturers
- National Ass'n of Soil Conservation Districts
- National Audubon Society
- National Farmers Union
- National Grange
- National Parks Association
- National Reclamation Association
- National Wildlife Federation
- Nature Conservancy
- Outdoor Writers Association of America
- Society of American Foresters
- Soil Conservation Society of America
- Sport Fishing Institute
- Wilderness Society
- Wildlife Management Institute
- Wildlife Society

ARIZONA LOCAL COMMITTEE

- Dr. Andrew L. McComb, Chairman, University of Arizona
- Joseph Arnold, Arizona State Land Office
- Robert V. Boyle, U. S. Soil Conservation Service
- William Carr, Pack Forestry Foundation
- Mrs. Hoyd Deckard, Arizona Federation of Women's Clubs
- Hon. Lewis H. Douglas, Valley National Bank, Tucson
- Mrs. H. Earle, Arizona Federation of Garden Clubs
- C. Edgar Goyette, Pioneer Hotel, Tucson
- Hal. Gras, Arizona-Sonora Desert Museum
- Dr. Reuben G. Gustavson, University of Arizona
- Mrs. C. R. Hensing, Arizona Federation of Women's Clubs
- Joseph Wood Krutch, Tucson
- Marvin Morrison, Arizona Farm Bureau Federation
- Dr. W. G. McGinnies, University of Arizona
- C. M. Palmer, Jr., Tombstone
- E. I. Rowland, Bureau of Land Management
- Robert J. Smith, Arizona Game and Fish Commission
- Dr. Lyle K. Sows, University of Arizona
- Jack Weadock, Arizona Daily Star, Tucson
- Mrs. Mabel Weadock, Tucson
- Norman Weeden, U. S. Forest Service
- William H. Woodin, III, Arizona-Sonora Desert Museum

PRESS ROOM

Rodeo Room on Mezzanine Floor
Daniel A. Poole, in charge

TICKET INFORMATION

Registration	\$ 5.00	
Luncheon, Monday, April 17	2.50	
Session II, Monday, April 17	7.00	
Includes all transportation and visit to "Water Street U.S.A." at the Arizona- Sonora Desert Museum; Reception, Chuck Wagon Dinner, Admission, and Entertainment at "Old Tucson"		
Luncheon, Tuesday, April 18	2.50	
Mt. Lemmon Bus Tour	2.00	
Reception and Annual Dinner	6.00	
Luncheon, Wednesday, April 19	2.50	
Total when purchased individually \$27.50		
Special combined price for all tickets .. \$25.00		
Post-Congress Tour of Walnut Gulch Watershed, Thursday, April 20		\$ 3.00

Eighth National Watershed Congress

709 Wire Building • Washington 5, D. C.

8th NATIONAL
WATERSHED CONGRESS

RAMADA INN

TUCSON, ARIZONA

APRIL 17, 18, AND 19, 1961



THEME: COUNT DOWN ON WATER

PROGRAM

374
e
ting

CONDITIONS UPON WHICH THIS COLLECTING PERMIT IS ISSUED

It is the intention of the National Park Service to further scientific research within the areas administered by it, and to cooperate with technical workers to the fullest extent compatible with its charge to preserve all species of flora and fauna and all geologic material in a natural state, insofar as is possible.

1. This permit applies only to animal life, plants, rocks and minerals. Archeological and paleontological materials may not be collected under this permit.
2. The collections shall be used for scientific or educational purposes only; shall be dedicated to public benefit, and shall not be used for commercial profit.
3. All collecting must be done away from roads, trails, and developed areas, unless such localities are specified in the permit. The collecting shall be conducted in such a manner as not to attract attention or to cause damage to the environment. Because of the scarcity or importance of some specimens, Service officials may designate the kind, number and sizes of specimens which may be collected, and any other restrictions necessary to the preservation of the area.
4. The National Park Service reserves the right, in the interest of science, to designate the depository of all specimens removed from a national park or monument and to approve or restrict transfers of specimens between depositories. The National Park Service also reserves the right to designate the U. S. National Museum as the depository of any type specimen removed from a national park or monument, after the collector has made necessary studies and published the results of his research thereon.
5. The Superintendent may require the permittee to furnish an inventory and locality description of any or all specimens proposed to be collected before they are removed and, after the collection is assembled, to submit it for examination.
6. Use or Disposition of Preserved Specimens: The collected specimens shall be deposited in a permanent public museum or in the exhibit, study or type collections of scientific or educational institutions. They must be suitably recorded in a permanent file and must be available to the public.

35548

egeneration project
National Research Council
Washington 25, DC, U.S.A.

Eighth National Watershed Congress
709 Wire Building • Washington 5, D. C.

P.S. If you send a cable you can use our Tucson Tel. number: East 6,9607. That helps get it to use quickly. Address it to me as MRS. WATSON SMITH.

WCS

P.S. If you send a cable you can use our Tucson Tel. number: East 6-9607. That helps get it to use quickly. Address it to me as Mrs. WATSON SMITH.

CLASS B

COLLECTING PERMIT
Restrictions Appearing on the Back Permission

United States Department of the Interior
National Park Service

In Accordance with the Conditions and Restrictions is Granted

Name of Collector F. Raymond Fosberg	Area Hawaii National Park	Date Issued March 27, 1961
To Collect the Following Specimens Plants, including one flower only from Hibiscalophus		Expiration Date April 10, 1961
Locality of Collecting Limited to Hawaii National Park		
Special Conditions or Restrictions Do not collect along trailside or roadside		
Approved (Signature) <i>Fred. Johnston</i>	Title Superintendent	

Two classes of collecting may be conducted under this permit:
 Class A - That required for public exhibits and for research undertaken by persons who can establish their connection with public museums or other scientific or educational institutions. Specimens collected may be insects (Hexapoda), spiders (Araneida), plants, rocks, or minerals, as designated in the permit.
 Class B - That undertaken by Federal employees only for scientific or educational purposes. Specimens collected may be plants, rocks, minerals or animal life as designated in the permit.
 The collecting of endangered or vanishing species of animals, if permitted at all, will be allowed only where the required approval has been obtained from the Director of the National Park Service

THIS PERMIT MUST BE CARRIED AT ALL TIMES WHILE COLLECTING. SEE REVERSE FOR CONDITIONS AND RESTRICTIONS.

F.A. Fosberg
Book # 59
begins with 41374
(nos. 41500-599 are in book 58 starting p. 52)



Standard
Miniature Blank
Book

No. 665 9 1/2 x 6 120 Pages Units



15658
62297

Vegetation Project
c/o National Research Council
Washington 25, D.C. USA.

P.S. If you send a cable you can use our Tucson Tel. number: East 6-9601. That helps get it to use quickly. Address it to me as Mrs. Watson Smith.

United States Department of the Interior
National Park Service

COLLECTING PERMIT

CLASS A B

In Accordance with the Conditions and Restrictions Appearing on the Back Permission is Granted

Name of Collector _____ Date Issued March 27, 1961

Area Hawaii National Park

To _____

P. _____

Loc. _____

F. _____

Spec. _____

D. _____

Appr. _____

By _____

Expiration Date April 10, 1961

Superintendent

undertaken by persons who can establish their educational institutions. Specimens collected plants, rocks, or minerals, as designated in the scientific or educational purposes. Specimens life as designated in the permit. if permitted at all, will be allowed only where the National Park Service

SEE REVERSE FOR CONDITIONS AND RESTRICTIONS.


THIS

F.A. Fosberg

Book # 59

begins with 41374

(Nos. 41500-599 are in book 58 starting p. 52)



Standard
Miniature Blank
Book

- No. 665 9 1/2 x 6 120 Pages Units
- No. 667 9 1/2 x 6 200 Pages Units
- No. 668 9 1/2 x 6 300 Pages Units

Made in the Following Rulings

- Journals, Day or Cash Books, Double S and Cts.
- S. E. Ledgers, S and Cts.
- Double Entry Ledgers
- Records with Margin Line

When ordering give Number and Ruling desired

Made in U. S. A.

A BOORUM & PEASE PRODUCT

Pacific Vegetation Project
c/o National Research Council
Washington 25, D.C. USA.

F. A. Fosberg

Book # 59

begin with 41374

(Nos. 41500-599 are
in book 58 starting
p. 52)



Standard
Miniature Blank
Book

No. 665 9½ x 6 120 Pages Units

No. 667 9½ x 6 200 Pages Units

No. 668 9½ x 6 300 Pages Units

Made in the Following Bindings

Journals, Day or Cash Books, Double S and Cts.

S. E. Ledgers, S and Cts.

Double Entry Ledgers

Records with Margin Line

When ordering give number and ruling desired

Made in U. S. A.

A BOORUM & PEASE PRODUCT

Pacific Vegetation Project
c/o National Research Council
Washington 25, D.C. U.S.A.

1960

Johnston Island

1

Oct 17 Visited at night - Terminal.

Thespesia populnea - planted
in front of terminal, to 3 m. tall.*Terminalia catappa* - 1 planted
in Terminal area, 2 m.*Cocos nucifera* - planted
trunks less than 1 m.*Chloris inflata* - abundant 7*Cyncha bonariensis* - abundant*Pluchea indica* - no seen.*Pluchea odorata* - " "*Cenchrus echinatus* - common*Cleusine indica* - occasional*Cyperus rotundus*

Kwajalein

ask for copy of report from
the station

Mr. Clegg - Preventive Medicine

Unit 6, Pearl Harbor.

Behind Dental Dispensary in P.H.

turn left, second turn to right

after entrance

Mr. Newsom, Hospital Administrator

says there are very few black
widow spiders here now

District Public Works Office

Fourteenth Naval District

memo to Commanding Officer

Pacific Missile Range Facility, Kwajalein

no. 3755 dated 20 May 1960

signed N.W. Martinson

3 spp. subterranean termites
1 dry wood termite*Formica purchasi* Mayh.*Chrysomya megacephala**Blattella germanica* (L.)*Periplaneta americana* (L.)*Rhodnius cardinalis*rats
mice

scabs

fly

termite

beetle
to control
scabs

1960 Johnston I. - Kwajalein Atoll

- Oct. 17 - Johnston Island
 41374 *Cyperus rotundus* L.
 locally forming patches
 around Terminal area.
- Oct. 19 - Kwajalein Islet
 on ~~coral~~ filled coral soil
 75 *Psilotum nudum* (L.)
 occasional between
 cobbles bordering walks
 around chapel.
- 76 *Hedyotis corymbosa* (L.) Lam.
 abundant, forming
 lawn in places near
 Terminal
- 77 *Eleocharis geniculata* R. & S.
 abundant very locally in
 marshy ~~open~~ spots
- 78 *Pluchea indica* & *odorata*
 very common locally with
 both parents in open flats
 and roadsides.

Oct. 21 - Kabbenbock Islet, Jaluit

- 79 *Hibiscus tiliaceus* L.
 very local, probably planted
 on coral sand and small gravel
- 80 fungus
 81 fungus
 on dead coconut log
- 82 alga
 abundant on shores of strongly
 brackish pond

41378a *Cynodon dactylon* ✓
 in meadow, ^{near} nesting nest.

prostrate. flowers
 very pale pink.

rather low, compared
 with both parents,
 stems tending to
 branch corymbosely;
 flowers mauve.

small twisted tree, 5 m. tall;
 leaves rather chlorotic;
 flowers bright yellow
 with maroon center.

Oct. 19 - Kwajalein Islet
 General appearance
 more luxuriant than I
 have seen it previously.
 Coconut trees bearing,
 around old Admin Bldg. site.

Planted or Cult. plants (a.w. = also seen wild)

Hymenocallis littoralis

Thespesia populnea

Pseuderanthemum carmenthonii

var. *atropurpureum*

Coccoloba uvifera

Hibiscus rosa-sinensis (a hybrid)

Nerium sp.

Pandanus tectarius

Cocos nucifera

Casuarina equisetifolia

Cinnamomum asiaticum

Lathyrus roseus

(a.w.) *Sponsea pas-caprae* var. *brasiliana*

(a.w.) *Tournefortia argentea*

Terminalia catappa

Ochroma oppositifolia

Pedilanthus tithymaloides

Vitex trifolia

Cordia alliodora

Clerodendrum unguiculatum

Native plants not planted:

Vigna marina

Scaevola sericea

Fimbristylis cymosa

Triumfetta procumbens

Wedelia biflora

Cassipouira bilobata

*Lepturus
reheus*

Weeds (* common at least locally)

* *Desmodium canum*

* *Hymenocallis littoralis*

* *Phyllanthus amarus*

* *Euphorbia prostrata*

* *Euphorbia thymifolia*

* *Cleome indica*

* *Euphorbia hirta*

* *Zinnia javanica*

Cyperus rotundus

Paspalum distichum

* *Paspalum longipes*

* *Cyperus compressus*

* *Tridax procumbens*

Cenchrus echinatus

* *Eclipta alba*

* *Stachytarpheta jamaicensis*

* *Hedyotis corymbosa*

* *Veronica cinerea*

Dactyloctenium aegyptium

* *Chrysopsis aciculatus*

* *Heliotropium ovalifolium*

var. *depressum*

Portulaca oleracea

Cynodon dactylon

* *Cyperus* or *Pluchea indica*

* *Phipalis angulatus*

Coryza canadensis

* *Bidens radiata*

* *Pluchea odorata*

* *Centella asiatica*

Euphorbia cyathophora

1960 Marshall Is.

On airstrip in sun in afternoon.
(2:50 pm.) 2 curlews (probably
whimbills, from their note), a
number of golden plovers, and
a considerable flock of turnstones.
Golden plovers common generally.

Old nursery in former
terminal - now being used
otherwise. Only a *Euphorbia*
pulcherrima bush, a clump
of tiny *Alocasia*, and a
vine of *Epipremna* *pinnata* (?)
climbing on a coconut tree
left of nursery stock.

Physalis *angulata* abundant.
A patch of *Ipomoea* *batatas*
cultivated in old *patio*
area.

Generally, the weeds that
were previously ~~not~~ found
only in the old terminal area
have spread throughout the
island e.g. *Bidens* *radiata*,
Cyperus *compressus*, *Paspalum*
longipes, *Heliotropium* *ovalifolium*,
etc. Some, such as *Coryza*
canadensis, *Pluchea* *indica*,
P. odorata, *Hedyotis* *corymbosa*,
& *Chrysopogon* *aciculatus*
have remained localized,
though ~~some~~ most of these
are very abundant locally.
The hybrid between *Pluchea*
indica and *P. odorata* has

Kwajalein

become very common
between the old and new
terminals, some variation
evident, but not much.

Few of the ornamentals
from the old nursery seem
to have persisted in the planting
around the island. *Hymenocallis*
is abundant. *Vitis* and
Pedilanthus somewhat so.

Oct. 20 - flight Kwajalein -
Majuro.

Passed Nanna on right.
Most islets are on the east side.
Passed Jabwoot - a very tiny
parcable, too far away to
see clearly. *Ailanthus* also.

Marshall Is.

Reef north from Majuro I. has a ridge exposed at low tide about $\frac{1}{4}$ back from seaward side. Small coral patches well scattered eastward in lagoon, ~~at least on south side~~ more numerous eastward, esp. on north side. Large patch with sand bar at low tide, about $\frac{3}{4}$ way east on north side where north reef begins in north reef with many islets and cross patches of exposed rock at low tide. Island on south side almost continuous.

Majuro - Jaluit flight. Reef flat very narrow along south coast of Majuro. No very appreciable algal ridge. Along south coast the reef flat is rather narrow. There seems to be an elongate pond occupying the place of the inlets in the intervals between islets on the south reef.

Circled Jaluit clockwise from n. of Mejatts ~~from s. of Mejatts~~ inside.

Bars along lagoon reef flat

Majuro - Jaluit

close to lagoon very common scattered rubble but no bars on outer reef flats. Surge channels all around, but very little development of algal ridge. Very striking bifurcating channels inside both passes. (photos beginning slightly after we first crossed reef mostly along east side from Mejatts to South Point, a few on south side, Luch and Lijeron Islets, some of submerged coconut trunks on east side at north and near Mejatts.

Marshall Is.

Oct. 20 - Jabwon, south of
old radio station.

Almost entire denuded area
has become covered by a mat
of *Wedelia*, *Vigna*, and
Sporobolus, extending almost
to the low cobble ridge which
is the ridge that was on the
reef flat. The *Sporobolus*
has sent a thin network
of strands many ^{meters} m. long
out towards the ~~the~~ cobble
ridge. The denuded flat
just back of the ridge
is only very thinly covered,
in places still bare.

Low scattered *Tournefortia*
and *Scaevola* locally
on this area. This mat
extends to perhaps 200 m.
beyond the old Weather tower.
Beyond this there is little
herbaceous growth, except
a few small patches of
Trumpetia and *Canavalia*.
There are scattered small
Tournefortia and *Scaevola*,
and abundant clumps
of sprouts from the old *Pennisetum*
stumps, these and a few small
Lepturus patches ~~with~~ roots
covered by sea water at high
tide. Many *Pandanus*
seedlings.

Jaluit Atoll

Between the Japanese Weather
Tower and Hydreytown there
is not a single standing
coconut tree or stump, dead
or alive, and only two or three
dead *Pandanus* snags, no
living ones.

Rough platform remnants
about midway between
Jabwon and Hydreytown
are not very much
indurated. ~~The~~ sample 11
from about 1 m. above
high, high tide level.
This layer is being
undercut in various
places, especially on lagoon
side, where looser layer
~~is~~ between two indurated
ones is exposed.

Very little loose material
on this part of land strip.
Here, for some hundreds
of yards, the ~~reef~~ ridge
that was on the reef has
vanished, at least from
sight at ~~the~~ high tide,
and it is not obvious
where the material has
gone. The vegetation is
of sparsely and irregularly
distributed *Tournefortia*
bushes to 1 m. tall, a few
small *Scaevola*, and a

1960

Marshall Is.

few dense clumps of *Pemphis*
sprouts along lagoon edge.

A few seedlings -

Scaevola, *Tournefortia*,
Sporobolus pes-caprae (1).

Two wandering tattlers on
lagoon reef.

Oct. 21, Jabuon -

Three muddy turnstones on
lagoon beach.

Polypodium scolopendria
Euphorbia hirta common
in *Wedelia* mat on
loose boulder deposit
along lagoon near
old radio station.

Terminalia catappa
trees here are growing
vigorously, there
than a sort of beaten
appearance, with
some dead branches, not
showing any effects
of typhoon.

A few *Pandanus*
trees alive and normal,
a few rather broken but
still alive. Some old snags.
Clump of *Casuarina* seem
in very good shape.

Coconuts planted generally
in pits. Mostly very
healthy, some are yellowish.

Jabuit

Kabbenbock

Oct. 21 - ~~British~~ 9

south fork of reef, in
of very coarse broken
coral, with locally some
bars of finer gravelly but
sharp light colored material
piled on top.

Islet said to have been
two islets, joined by an early
typhoon.

Inner end of islet
surrounded by coarse
beach rock, some finer.
Small beach ridge of coral
gravel.

7 trees seen.

Tournefortia

Pandanus

Scaevola

Cordia

Marringtonia asiatica (seedling)

Mouroua citrifolia

Quettarda

Hernandia

Terminalia

Hibiscus tiliaceus

Herbs

Crinum

Wedelia

Fimbristylis

Sporobolus tuba

Hedyotis biflora

Polypodium scolopendria

Asplenium nidus

Canavalia microcarpa

Vegetation

Signa marina

Lepturus

Tacca

Euphorbia hirta

1960 Marshall Is.

- 1 little-thighed curlew
- 2 fairy terns
- 3 common noddies
- 2 tattlers

2 reef herons, one speckled, one white with ^{small} ^{beak} ^{shoulder} ^{patches}

On seaward side is a small embayment that has been dammed off probably by the typhoon, by a ridge of gravel, enclosing a pond. water not quite salty as sea water, abundant green algae, some blue green.

cordia. Tipped over fernandis and terminalia trees growing vigorously. Healthy upright saplings of Terminalia.

Front reef facing passage has a broad relatively smooth fine grained planation surface, perhaps 100 m. wide, sloping off rapidly seaward, with active coral growth for 5-6 m. down. Then debris to flat sandy (?) bottom of pass (see Knudson). Landward a very flat erosion ramp up to 100 m. wide (in embayment) much narrower elsewhere, fairly rough, peeling off at edges (see photos). a few and in layers.

remnants of beach-rock to south.

Platform here at about high tide level. Scattered medium sized boulders on reef flat and ramps. up to 4.5 m. diam. irregular.

+ some fair sized areas of old surface that seems to be an old growth surface of ~~the~~ Acropora.

Scattered boulders of vesicular lava from ballast of wrecked ship

On inner shore there is an extensive low-tide level plate with a rather narrow platform, perhaps 8-10 m. wide, protruding from ~~the~~ islet, with narrow erosion ramp, peeling off in 2 layers. Undercutting + collapse (see photo) (see photo of platform).

Pieces of ^{seems} ^{somewhat} vesicular black volcanic rock up to 20-25 cm. long scattered on boulder beach on inner side, apparently carried around from the wreck on the outer ~~to~~ reef, abundant on outer beach.

Very few coconut trees left on this islet, a few of those lying down have turned up at tide

1960 Marshall Is.

From lagoon about 3 standing coconut trees visible in vicinity of Hydreytown, mostly around tanks.

Jabwon -

1 whimberl on seaward side.

Hemigraphis reptans around old masonry foundations, barely holding on.

Hymenocallis littoralis very common very luxuriant, covered by sea water at high tide.

Bananas planted abundantly and growing luxuriantly around old ruins.

Breadfruit - vigorous sprouts from old uprooted tree, small upright tree in very good shape.

Cyperus thyllingia locally abundant but dwarfed.

Ocimum sanctum,
Catharanthus roseus, *Mirabilis* and yellow hybrid *Canna* planted in front of houses doing well.

A few patches of sweet potatoes in sheltered spots doing very well.

Jaluit

Major 1922

Oct. 2 - Mejiro (Elizabeth)
Aspect in village not strikingly different from pro. typhoon appearance - trees and cultivated plants generally young.

Plants seen:

- pl. o *Plumeria rubra*
pl. o *Polyscias scutellaria*
c *Premna obtusifolia*
a *Pandanus tectorius*
pl. o *Nerium indicum* (sprouting from stump)
pl. c *Musa sapientum*
pl. a *Cocos nucifera*
a *Fimbristylis cymosa*
pl. c *Artocarpus altilis*
c *Morinda citrifolia*
a *Polypodium scolopendria*
a *Lepturus repens*
pl. a *Nephrolepis acutifolia*
pl. a *Prunus gymnorhiza*
c *Canna papaya*
pl. c *Cucurbita pepo* (round fruit, spotted leaves)
pl. o *Xanthosoma sagittifolium* (chlored)
o *Tacca leontopetaloides*
pl. c *Cleome indica*
pl. a *Triumfetta procumbens*
c *Pteris tripartita*
pl. c *Conchocarpus echinatus*
pl. a *Eragrostis amabilis* (in paths)
pl. *Alocasia macrorrhiza* (planted)
(leaves very erect. The angle with blade.)
a *Vigna marina*
pl. o *Codiaeum variegatum* (to n. 1100)

18

1960 Marshall Is.

Oct. 21 - Jabwor

41383

Stachytarpheta urticifolia
one plant in open herb flat

84

Hemigraphis leptans (Forst.) And.
rare around old masonry
ruins.

Oct. 22 - Mejinilo (Elisabeth) I.

85

algae
~~near~~ just below high
tide level in tidal pond.

86

(algae)
epiphytic on *Pemphis* trunk
near high tide level around
tidal pond.

87

moss

88

moss

same

Oct. 22 - Jaluit Islet

89

(fungus) *Dicodonia elegans* Forst. 215
on rotting coconut trunk

90

Summitzera littorea
locally abundant in
swamp.

91

Sonneratia alba
locally common in swamp

92

Vittaria incurvata Cav.
rare, epiphytic on tree
trunk in moist forest.

Jaluit Atoll

19

branched herb to 0.8 m. tall,
flowers deep purple.
prostrate; chlorotic.tree 4 m. tall (others taller);
leaves sub-fleshy to leathery;
flowers scarlet "kimeem"
tree 6 m. tall, spreading
(others erect, to 10-12 m. tall);
petals linear, white,
stamens white "pulabl"
"sweet nectar from flower
eaten by Marshallese.

1960 Marshall Is -

Several young coconut trees (nuts from Majuro) the leaves, after the first 204, not completely opening up - pinnae adhering at margin, separating along midrib. 3 seen in immediate area of village, many other seedlings normal.

In coconut groves the general ground cover - mostly *Lepturus*, is complete.

Much *Fimbristylis*
Patches of *Nephrrolepis hirsutula*, scattered
Pteris tripartita + clumps of *Polypodium*, patches of *Triumfetta*

Many young breadfruit trees planted, some volunteers, about 2.5 m. tall, expected to produce fruit in from 6 to 11 years.

Many small bushes of *Mouroua*, *Premna*, scattered in coconut grove. Mostly appearing to be from old cut stubs, possibly from clearing for coconut planting. Bananas bearing, look healthy except occasional slight chlorosis.

Plants seen (etc.)

- ae *Pseuderanthemum carmichaelii*
- lc *Digitaria pruriens* v. *microbachne*
- ea *Cassytha filiformis*
- ea *Wedelia biflora*
- eo *Euphorbia charnissii*
- ea *Fleurya ruderalis*
- l *Portulaca oleracea*
- eo *Guettarda speciosa*
- ec *Scaevola sericea*
- ~ *Allophylus timoriensis*
- eo *Cordia subcordata*
- ~ *Terminalia samoensis*
- ~ *Asplenium nidus*
- ~ *Calophyllum inophyllum*
- ~ *Procris pedunculata*
- l *Pamphis acidula*
- h *Pipturus argenteus*
- l *Peperomia forapensis*

Patches of pumpkins planted, also small patch of *Alocasia*. Some *Pandanus* planted for food. Volunteer *Pandanus* used for hats + mats.

On ^{side} end of islet near passage to e. *Polypodium*, *Cassytha*, *Triumfetta*, *Nephrrolepis* abundant in ground cover. Some *Euphorbia*, *Wedelia*, *Digitaria*, *Fleurya*, *Fimbristylis*, aspect drier.

~~*Wedelia* generally not abundant, as *Fleurya* in rocky places~~ ^{except on the passage side}

photos in open area back of lake, filled with *Wedelia*. also an area of young coconuts and across lake, all from same spot. (low tide)

Patch of *Pemphis* around edges of lake.

Algae not conspicuous in lake as before, but still present.

Seedlings of *Pemphis* and *Bruguiera* abundant around margins of lake, none in lake bottom, though it is almost dry at low tide. (photos)

Wedelia is abundant and forms blankets in open areas toward seaward side, especially near east passage.

In semi-open area, with coconut trees with trunks 1-1.5 m. tall, *Polypodium lepturus* and *Cassytha* most abundant. *Cassytha* thicket in open spots. seedling *Morinda* common here.

Cordia trees ~~are~~ occasional along beach ridge, not inland, no seedlings. *Guettarda* not common inland from sea passage.

Mounted
Salpene
Siphon

There are still many coconut trunks lying on ground, some rather well rotted, others still firm, but in general rather well cleared away, where they formed an obstruction.

Back of *Pemphis* zone back of lake in open area generally covered by *Wedelia* are coppice-like clumps of seedling *Guettarda*, possibly cut off once but abundantly sprouting, about 1.5-2 m. tall. (photos)

Bottom of lake hard, pan-like, level, mainly, but with pits near center with water at low tide. ~~shrimp~~

No mud except very near edges around mangrove and *Pemphis* patches. (sample 12)

On old gnarled *Pemphis* trunks epiphytic algae, mosses. *Nephrolepis acutifolia*, *Coccoloba*, *Polypodium*. On older *Bruguiera* epiphytic *Nephrolepis acutifolia*, large clumps.

photos
Palaeomon
(Palaeomon)
delibis
Dome

1960 Marshall Is.

On n.w. corner, even in rather open areas, on Wedelia, Polypodium & Nephrolepis & seedlings. Pandanus forms principal ground cover, sparse & 3m. tall with Trimmetha, Fimbristylis, Lepturus, Fleurya, Polypodium, Nephrolepis, seedlings. Tournefortia on beach ridge (sharp gravel). Scaevola seedlings just above reef flat.

In areas with semi-closed cover of coconuts of various sizes, a luxuriant tall green cover of Polypodium and Nephrolepis about 1 m. tall. Large black streak seen.

On ~~the~~ n.w. corner is a patch of thicket with tall coconut trees, Coppice like Pipturus to 4 m. considerable shrubby Altophyllus, much Asplenium and Probris, Peperomia, etc. growing on deposit of large sharp boulders. Ferns luxuriant. In some shrubby Datisca, Ficus, Morinda. This area is a boulder tract perhaps

1 m. above general level of island.

Just inside this is park-like coconut grove, Lepturus forming dense ground cover, seedling Datisca, clumps of Nephrolepis on fallen coconut butts, surrounded by thickets of Pipturus, Morinda, Pemphis, Pandanus seedlings. Small Bruguiera depression, this filled with rotting coconut husks, probably left by typhoon. Pandanus trunks pretty well rotted. Coconut trunks mostly still rather firm, but bark coming off.

Southeast corner rather open several well formed young breadfruit, a few ragged Pandanus, one or two bare old breadfruit trees. scattered coconuts. general dense ground cover of Vigna, this rather chlorotic. (photos). One small patch of Wedelia (dark green).

Plants seen (det.)

- lc *Intsia bijuga*
 lc *Thunaea involuta*
 sl. o *Cinnamomum procerum*
 pl. o *Mirabilis jalapa*
 pl. n *Hippocrepium pumilum*
 pl. n *Catharanthus roseus*
 pl. n *Tapeia*
 a *Physalis angulata*
 pl. n *Zephyranthes rosea*
Asclepias curassavica (in leaf)
Polycaea butirosa (in leaf)

Coconut planting has been fairly general, regardless of whether there were trees not damaged by typhoon, except for very young ones.

Several turnstones on bottom of pond.

2 reef herons one white, one almost white.

Flock of white-capped noddies in lagoon
 several white terns inland

Smaller islets on south reef, mostly have a tangle of scrub with a few coconut trees, none of these very tall. Much of tangle, at least on lagoon side, seem to be *Scaevola*.

Large islands mostly well covered by coconuts, some leaning at odd angles, but mostly looking rather normal.

Considerable sand on lagoon sides.

Bakan - islet nearest Jaluit I. seems to be 2 small islets joined together - rather few coconuts except on west end. There on east end rising through irregular thickets of small trees and shrubs.

Reef between islets very wide, inner slope rather steep, covered with rubble debris. Opposite islets more sandy, gently sloping, much live coral in shallower parts.

Wide shallow area opposite Jaluit I.

Oct. 22 - Jaluit I. e. end.

Plants seen Littorina
Pandanus tectorius
Coccoloba ovifera
Plumera rubra
Carica papaya
Lepturus repens
Musa sapientum
Cucurbita
Mirabilis jalapa
Hymenocallis littoralis
Cuminum asiaticum
Artocarpus altilis
Zephyranthes rosea
Phyllanthus amarus
Gomphrena globosa
Euphorbia hirta
Eleusine indica
Huaca involuta
Digitalis purpurea
 var. *microbachne*
Cenchrus echinatus
Wedelia biflora
Alocasia macrorhiza
Polypodium scolopendria
Tacca sp. leucopetaloides
Nephrrolepis bisentata
Vigna marina
Pemphis acidula
Clerodendrum inerme
Trinostylis cymosa
Trinostylis procumbens
Scaevola sericea

Tournefortia argentea
Bougainvillea gymnantha
Pipturus argenteus
Guettarda speciosa
Flourensia ruderalis
Asplenium nidus
Nephrrolepis acutifolia
Vittaria elongata
Pisonia grandis
Allophylus timoriensis
Sonneratia littorea "hincen"
Sonneratia alba "palab"
Mimosa catapappa
Vigna marina
Cassipouira filiformis
Centella asiatica
Blechnum pyramidatum
Cyperus javanicus
Crinum procerum
Calophyllum inophyllum
Portulaca oleracea
Terminalia catappa

Several golden plovers inland
 (whimbrel in swamp)
 white tern in swamp

East of village, a low completely bare irregular area showing signs of flowing water, drain from swamp?.

West of this is an area of an area where all the coconut trees have been cut down - even apparently healthy rather young ones - in order to let in sun for the newly planted young ones. (photos to be taken)
A few tall Pandanus left standing, surrounded on one side by Pemphis forest. The Pemphis forest is really a tidal swamp irregular, mostly about 5 m. tall, no undergrowth at all. Bare coarse sand as in adjacent open bay flats.

In these some *Burquiera* but most of this was apparently destroyed by the typhoon. (photos to be taken)

West. East of this a rather extensive ~~area~~ strip of clumps of Pemphis ~~and~~ sprouts about 2-3 m. tall, open. Then a large area that was *Burquiera* swamp, but all old trees

removed by typhoon - root and kernel systems remaining. Much of this now covered by young *Burquiera* saplings 1-3 m. tall! (photos to be taken)

Between this and beach a sheet of coarse gravel perhaps 200 m. wide with a few standing coconuts, many fallen ones. Open stand of young clumps of *Tournefortia* and *Peacock*, about 1 m. tall or less.

Planted to coconuts: some patches of *Lepturus*, some *Trumbetta* inland from beach & few snats of *Wedelia*. (photos - to be taken)

This gravel sheet swept into edge of mangroved swamp, margin lobed, about 0.5 m. thick bed.

Trees here have fallen either east or west or generally ~~that~~ ^{both} of these directions.

Considerable acreage of *Burquiera* saplings, a few old trees left in s.e. corner of swamp.

Wedelia is abundant toward s.e. corner of islet, outside swamp.

As pest here generally wides, more luxuriant
Aplenum large,
Pipturus common,
Euphytes common.

On south side of swamp
Sunnitiera is common
 and a tree of *Tournefortia*.
 Much of this swamp
 is hard bottomed,
 some covered by small
 broken coral debris.

In this area *Sunnitiera*
 is locally dominant
 in swamp.

Not much damage
 here by typhoon.

This area of swamps,
 tidal flats, localized
 higher ground extends
 westward through middle
 of islet. Locally *Tournefortia*
 is abundant, with some
Bougainvillea, *Sunnitiera*,
Pemphis. Locally *Bougainvillea*
 or *Pemphis* more abundant.
Pemphis usually on slight
 higher ground.

One area between swam
 and lagoon almost
 treeless, grassy.

Back of lagoon ridge in
 places are marshes,
 mostly dominated by
Cyperus javanicus.

Pandanus not at
 all abundant on this
 islet.

Cong Islet.

On arcuate gravel bar
 along lagoon shore, interrupted
 a few scattered ~~see~~ young
Scaevola, *Tournefortia*, 2
Cocos seedlings.

One white reed here.

Pemphis abundant
 along lagoon shore -
 clumps of sprouts.

Oct. 23 Enybor I.

Plants seen:

- lc Antrocarpus altilis
- la Cocos nucifera
- c Morinda citrifolia
- c Pandanus tectorius
- c Pipturus argenteus
- a Wedelia biflora
- la Vigna marina
- lc Asplenium nidus
- lc Physalis angulata
- la Peperomia acidula
- ~ Alocasia macrouriza
- o Polypodium scolopendria
- ~ Kalanchoe pinnata
- la Lepturus repens
- ~ Allophylus timoriensis
- ~ Pisonia grandis
- o Quettarda speciosa
- la Hibiscus tiliaceus
- lc Scaevola sericea
- ~ Calophyllum inophyllum
- lc Premna obtusifolium
- c Tournefortia argentea
- lc Fleurya ruderalis
- l Portulaca oleracea
- lc Phyllanthus
- l Cyperus javanicus ?
- lc Fimbristylis cymosa
- l Digitaria pruriens v. murd.
- ~ Cassytha filiformis
- ~ Tacca leontopetaloides
- lc Thuarea involuta
- ~ Cenchrus echinatus
- ~ Eleusine indica

Small flock of common noddies on old tree skeleton near wrecked pier and generally flying around passage end of islet.

Grove of breadfruit trees perhaps 20-30m dbh. Broken off part way up by typhoon - almost all have sent out vigorous sprouts from upper part of trunks and formed full-crowned well-formed trees, now flowering.

Ground cover generally Wedelia, locally Sigra. Has been cleared out some time ago. Some breadfruit trees nearer sea have not recovered, are completely dead.

Lined skinks abundant.

Wedelia seedlings in great abundance especially where there is decomposing trash and on rotting branches.

Pipturus sprouting vigorously from stumps where trees have been cut during clearing. Uncut trees commonly 5-8m tall. Many of sprouts chlorotic. These heavily infested with red mites. (coll.) Aedes abundant (coll.)

In the pass between Ruytor and Kabbenbock is a Y shaped reef with very long arms, the right one perhaps is more or less a number of very large boulders, and much large rubble.

A reef patch in the right hand channel about even with the inner end of the above branch of the Y shaped reef.

These all bare at low tide. ~~Some~~

A considerable number of large volcanic boulders from wreck on Kabbenbock here around landing, probably brought by people.

~~Some~~ Hibiscus

~~Some~~ ^{some} Hibiscus tiliaceus, around a small ponds, some not. Certain branches very chlorotic mostly normally green ^(chlorotic)

Most of the area of this islet has been almost completely cleared, except clumps of Pemphis, Tournefortia, occasional breadfruit tree. Covered mostly by a blanket of Wedelia.

One area has had the Wedelia chopped away, rolled up and burned recently. Here the gravelly dark gray soil is being colonized by millions of seedlings of Wedelia.

Many bomb craters filled with water, green with microscopic algae.

Several white terns flying around, mostly near Hibiscus thickets.

Common noddies by far the most frequent birds.

Many young coconut trees, from before typhoon, along pass and lagoon sides.

~~Hibiscus~~ The cleared area has been staked out for planting coconuts but very few have been planted.

On sand flat protruding into lagoon are seedlings of Scaevola, Tournefortia, Triumphetta, Pandanus, Wedelia, Lepturus, Vigna, Sponges pericarpia (chlorotic)

Plants seen (ctd)

- P.d. *Pseuderanthemum canuttii*
 l. *Hernandia sonora*
 a. *Sophora tomentosa*
 l. *Sporosa pec-capras*
Terminalia catappa
Caesalpinia ~~rostrata~~ bonduca
Musa sapientum
Sonneratia littorea

1 reef heron and 1 tattler seem to be
 companions, when one flies or alights, other do
 1 white, 1 white with black wing tip
 1 white heron, 1 blue, 1 ~~blue~~
 grey with white shoulder patch
 several wandering tattlers
 on seaward reef flat
 1 turnstone on lagoon beach

young coconut trees
 seem mostly decumbent
 at base, then erect.

In lagoon shore many
 small *Rattus exulans*
 running around in broad
 daylight, scarcely afraid
 of humans.

One narrow part of islet
 is a long masonry stone wall
 about 0.7 m. high
 running from place where
 islet narrows almost
 to first cluster of buildings.
 On seaward side gravel is
 piled against it part
 way, no vegetation.

In lagoon side a narrow
 reef platform strip with
 a slight accumulation
 of sand at lagoon edge.
 Then vegetation of *Sigra*,
 small *Panicum* wisps,
 small *Transepedia*
 bushes, small *Leucaena*,
 coconut seedlings, a few
Pandanus seedlings, some
Repturus, esp. on sandy place,
 but also on bare rock, *Triumfetta*
 A few larger coconut trees,
 a few *Hibiscus* seedlings,
 Much bare rock.

Beyond end of wall no veg. on
 bare eroded platform (except 1 small
Transepedia & *Urtica* clump of buildings).

Sporosa
pec-capras
 seedlings

lagoon
 reef flat

at end of sea wall
the platform rock is
eroded, a few scraps of
upper surface left the
indurated, below this
merely compacted - firm
but crumbles easily when
hammered. Black with
algae. Topmost parts of
this platform must
be 3 m. above l.t.

sample #13 - indurated flat
surface

sample #14 - softer material
below.

sample #15 - hard material
from rough surface of
seaward erosion ramp
cut in coarse platform breccia

ONEIDA

Oct. 23 Jabwon

Terminalia catappa
fairly well leaved out
or sprouts all over old
trunks and broken branches.
A few Pandanus with
big tufts of leaves on
branches that were not
broken. No sprouting
from broken branches.

Fairly extensive sweet
potato patches that
look pretty good
along main street
north of radio station.

#15 - Along path

- a *Euphorbia prostrata*
- a *Phyllanthus amarus*
- a *Veronica cinerea*
- a *Euphorbia hirta*
- a *Pilea microphylla*
- a *Dentella repens*
- a *Cyperus rotundus*
- a *Portulaca oleracea*
- a *Lynedrella nodiflora*
- a *Crotalaria incana*
- a *Distachytrapheta verticillifolia*
- a *Eleocharis pyramidatum*
- a *Cyperus pennatus*
- a *Wedelia biflora*
- a *Cragrostis amabilis*
- a *Leucaena glauca*
- a *Cyperus compressus*
- a *Eleusine indica*
- a *Hedyotis corymbosa*

Euphorbia prostrata
Phyllanthus amarus
Veronica cinerea
Euphorbia hirta
Pilea microphylla
Dentella repens
Cyperus rotundus
Portulaca oleracea
Lynedrella nodiflora
Crotalaria incana
Distachytrapheta verticillifolia
Eleocharis pyramidatum
Cyperus pennatus
Wedelia biflora
Cragrostis amabilis
Leucaena glauca
Cyperus compressus
Eleusine indica
Hedyotis corymbosa

Ornamentals

Lantana camara
Pseuderanthemum carmichaelii
 and var. *atropurpureum*
Zephyranthes rosea
Plumeria rubra
Hymenocallis littoralis
Mirabilis jalapa
Zinnia elegans
Catharanthus roseus
Gomphrena globosa
Quimium sanctum
Canna hybrid

Ageratum, very rare before,
 now common on distal
 part of island. *Cyperus*
rotundus, observed as a
 small patch in village,
 now scattered around
 trails faintly generally.

Crotalaria still very common
 but by no means as much as
 as before - largely crowded out
 by *Wedelia*. *Blechnum* very
 abundant, especially where
Wedelia has been cleaned out.

Terminalia, *Hernandia*,
Calophyllum, *Artocarpus*, *Ficus*,
 have recovered very well, filling
 out crowns with sprouts (except
Artocarpus when too damaged,
 then it died).

In old botanical garden
Epipremnum pinnatum
Cycas circinalis
Ficus elastica
Delonix regia
Erythrina variegata
Schinus molle
Scaevola taccada
Cassia occidentalis
Hibiscus tiliaceus
Calophyllum inophyllum
Lamanea saman
Rhoeo spathacea
 KNOX

Distal part of island
 almost completely cleared,
 covered by a clipped weedy
 growth of *Blechnum*, *Stachy-*
tarpheta, *Leucaena*, *Vigna*,
 etc. about 2-3 dm high. Weedy
 feathered trees - *Carica*,
Terminalia, *Artocarpus*,
Pipturus, *Pandanus*, *Musa*,
Hernandia,

2 whimbrels flew up
 near radio station
 heard a brush-tufted
curler.

Oct. 23 - Enybor Islet

- 41393 *Caesalpinia bonduca* (L.) Roxb.
2 single plant on narrowing
part of islet in deep tangled *Wedelia*
- 9 94 *Phyllanthus amarus* Schum. & Thonn.
abundant in recently
disturbed places
- 6 95 *Hervandia sonora* L.
occasional
- 2 96 *Pandanus* ~~sp.~~
common

Oct. 23 - Jabwor
in waste spots and
along paths

- 2 97 *Dentella repens*
abundant along paths
- 3 98 *Hedyotis corymbosa*
abundant along paths
- 2 99 *Euphorbia thymifolia* L.
very local in path
- 41400 *Cyberus bylingia*
2 very local along trail
- 2 01 *Digitaria pruriens* var. *microbachne*
very local in trail
- 1 02 *Euphorbia prostrata*
very common in trail
- 5 03 *Euphorbia glomerata* (Millsp.) Weibull
locally abundant
Oct 25 Long Islet bet Hydneton
and Lelud
- 1 03 *Hibiscus tiliaceus*
rare seedlings on bare gravel
- 1 04 seedling

young plant, three
vine-like branches 1-2 m.
long; sterile.sprouting fallen tree;
fruiting envelope inflated,
sifia not toothed at all.
tall tree; fruits on
ground.prostrate, forming a
mat, corolla pale dull blue,
lobes with white margins.
prostrate, forming a mat,
corolla white.
stems and leaves green,
flowers whitish.
heads white

(P. rot) Fish culms spreading.

whole plant purple.

erect

Oct. 24 - Long Islet, bet. Jabuon and Jaluit.

(brow. photo) In the area where the ridge of boulders on the seaward reef flat has disappeared, at low tide it becomes obvious that it has largely just spread out and become a rubble tract.

On lagoon reef flat there is a broad flat strip of gravel separated from the land-strip by 30 to 100 m on the surface of this blackened rather stable mass is a ridge of smaller material, obviously more worked, whiter, forming a sharp-topped bar ridge.

photos

This is more a less the situation, the length of this strip between Jabuon and Jaluit.

Seedling of *Calophyllum* deeply rooted, healthy, in bare gravel sheet.

Permpis forest - bare trees still standing, trunks

sometimes sprouting, clumps of sprout up to 1-1.5 m tall at bases (photos).

Just north of Sydneytown an area of low platform awash at high tide quite abraded. Inside this the lagoon bar is missing, outside, the gravel ridge on the reef flat is still mostly separated by 15 m. or so of water from the land strip. This ridge is at least 2-2.5 m. high in places, quite abrupt on inner side, imbricated slabs on outer side. (photos)

This outer ridge is lacking opposite Sydneytown but a sharp ridge is piled up on the outer edge of the land-strip just n. of Sydneytown.

Several wandering tattlers on outer reef flat.

On lagoon reef flat opposite a cut through the beccia platform is an ^{at least} inner gravel bar, inside the main one. (photo - hood.)

The end of the main outer bar here is just opposite the cut in the breccia platform and is rounded, imbricate seaward around the end, ~~to~~ and the opposite side of the cut is swept very clean, suggesting that the currents or waves come through this channel from a dominantly n.e. direction, exerting their force against the south side of the channel, and arranging the gravel slabs on the lobe of the ridge in imbricate fashion with less energy involved, thus not carrying this loose material along into the lagoon.

Channel ~~the~~ dug across land strip by Japanese south of Sydneytown in three layers, successively harder downward (photos - b.w.), samples 17, 19, 19 (downward).

To lagoon side of this is a circular debris bar, surrounding a pool. (photos), this immediately

lagoonward of shore, a broad reef flat ~~starts~~ lagoonward of this, then a well developed bar on lagoon edge of this. The flat well strewn with large boulders.

Seaward the boulder ridge is separated from the erosion ramp, and here, for a ~~great~~ distance of perhaps 100 m, loops out away from ~~the~~ shore and is higher than in either direction, where it is spread out and low.

South of here a very low strip, about 100 m. w., scoured off except for remnants of an old breccia platform. Lagoonward is a concentration of debris - a high crooked place in the debris ridge (photos - b.w.) a number in several directions. Seaward from this wash another loop of high ridge, again extending a bit south of beach in land strip. South to here on land strip, for Sydneytown vegetation, mostly a low irregular scrub of *Tournefortia* and *Scaevola*. Pamphion ^{some of lagoon} southward same but ^{very flat} much lower and thinner. (photo)

Enormous boulders
thrown upon lagoon slope
of lagoon ridge, up to
2 x 2 x 3 m. but irregular.

Many of them, this the
largest. Some are large
Millepora colonies, most
are very complex. A few
of these are erosion remnants
of a former platform.

This lagoon ridge,
too, is broad and thin
and lies on a planation
surface that is much
like an erosion ramp,
sloping slightly lagoonward
about 10 m from top of
seaward edge of this.

(Photos showing ^{or} remnant
on left, boulders on it.)

Just beyond, a breach
in landstrip, ^{showing}
that the reef ^(with a broad erosion ramp) flat continues
under the consolidated
debris ridge to lagoon
side. much denser
and harder than the
material lying on it.

In this dissected partially
consolidated material
there are old stumps of
Pemphis on lagoon side.
Young plants of Tournefortia
and Scaevola making

up an open scrub,
with ~~also~~ ~~some~~ small
mats of *Triumfetta*.
Seedlings of *Scaevola*,
Tournefortia, *Pemphis*,
Cocos, *Vigna*, *Hibiscus*,
Clitorea, *Barringtonia*
asiatica, *Pandanus* (in
loose debris).

Two white terns flying.

Pemphis, *Scaevola*, ^{*Laportea*} *Tournefortia*,
Cocos, *Fimbristylis* can
clearly grow where inun-
dated by very high tides.

More vegetation on
higher spots, also larger
Tournefortia and *Scaevola*
plants. Patches of *Triumfetta*.

Photos of broad erosion
ramp and rubble tracts
on seaward reef flat,
also of scrub on debris ridge
on landstrip beyond ^{most} concrete
building, toward warehouses.

On debris ridge seedlings
of *Guetarda*, ^{*Canavalia*} *Barringtonia*, *Cocos*.
Stump of *Tournefortia*
& *Scaevola* 1-2 m high.
Ipomoea tuba in it.
Young coconuts.

Sharp ridge near
lagoonward edge of
lagoon reef flat -
being colonized by *Pennisetum*,
Pennisetum, *Commersonia*,
Vigna, *Hibiscus tiliaceus*
Tournefortia seedlings.
Pandanus fruits thrown
up but not seen germinating

Wide place at warehouse
with open scrub of
Tournefortia, *Caeculia*,
Pipturus, a little
Allophylus, patches
of *Wedelia* occasional
Pennisetum + *Allophylus*.

some dense residual
scrub of ragged *Pandanus*,
Pisonia, *Guettarda*, *Tourne-*
fortia, Coconut.

This area seems to be
a large gravel sheet
laid down by the typhoon,
the residual vegetation
not quite buried. Most
of the rest representing
colonization. Surprising
thing is number of
bushes of all sizes of
Pipturus. Some of the
smallest ones seem to
be drying up, but mostly not

Here a sheet of very
rounded flat gravel
extends from the land strip
out onto the lagoon reef flat
with a lobed margin.

Much open ~~open~~ bare
gravel between bushes,
but some large mats
of *Wedelia*, *Vigna*,
Triumfetta, *Canavalia*
and *Ipomoea pes-caprae* (thin)
have small patches of
Fimbristylis.

Reef herons seen - 3 white,
1 blue, 2 white with some grey,
1 whimbrel.
1 reef heron chasing or playing
with a tattler.

Beyond warehouse we went
by boat. Generally the
vegetation is a strip of
~~open~~ and irregular
scrub, mostly *Tournefortia*,
on platform and debris,
with sparse *Pennisetum*
on denuded flats or erosion
ramp.

Standing coconuts very
few around warehouse,
even fewer down the land
strip, then becoming

slightly more abundant till we come to a wider place, where a considerable number are left standing. Here at this wide place, a high ~~ridge~~ ^{gravel ridge} along lagoon shore, and on beach of this huge coral blocks scattered.
2 whimbrels here (!?)

Here the gravel ridge has mostly moved onto the shore, making a typical boulder ridge, but to note it is still a bit separated.

On the flat is an open to closed scrub of *Tournefortia*, some *Scaevola*, some *Quettarda*, many stunted seedling crowns, incomplete groundcover of *Trunifetta*, some *Sporogon tuba*, some *Pandanus* seedlings, some *Canavalia*. ^{standing crowns}

Part of this that was ^{scoured} by typhoon is pretty well vegetated, with even some residual sprouting *Quettarda*, etc. Part covered by grand sheet has scattered bushes of *Tournefortia*.

Scaevola up to 2 m tall.

Lagoonward these become smaller, up to 0.5 m.

Pemphis seedlings and clumps of sprouts on hard rock near lagoon, some curved gravel bar near lagoon shore, being colonized by *Pemphis*,

Lepturus, *Tournefortia*,

Scaevola, *Pandanus* (s.)

Wedelia (n.s.), the plants up to 1 - 1.3 m tall.

Pandanus trunks on ground rotten.

Southward a similar scrub, with scattered standing coconut trees, becoming lower and more sparse in places, the *Pemphis* along the lagoon varying from dense to absent, always principally on bare erosion ramp or platform rock.

Many bare *Pandanus* trunks still standing.

Lagoon bar discontinuous, high in places.

One stretch of it has many great boulders strewn on its lagoon slope. Clearly from lagoon bottom.

Oct. 24 Jabon -
 photos (b+w) of trees along
 lagoon shore of Jabon.
 1 blue, gray, white reef beam
 on end of Jabon.

Oct. 24 - Enybon -
 photos - b+w, Kod. of
 recovered breadfruit
 trees on N. W. corner of islet.

Oct. 25 - crossing lagoon
 Jabon to Pinlep - one brown
 booby flying over lagoon.
 returning in evening & brown boobies.

Oct. 25 - Pinlep

Plants seen

- la Lepturus repens
- ec Cassytha filiformis
- e Centella asiatica 'manila'
- c Trionfetta procumbens
- c Thuarea involuta
- c Pandanus tectorius
- la Digitaria pruriens v. microbachne
- a Crocos naurera
- ec Carica papaya
- a Vigna marina
- la Phyllanthus amarus
- o Cerebrus echinatus
- ec Eleusine indica
- la Euphorbia hirta
- ec Eragrostis amabilis
- c Artocarpus altilis
- M Catharanthus roseus
- pl Cinnam procerum?
- M Musa sapientum
- a Wedelia biflora
- ec Tacca leontopetaloides
- a Pipturus argenteus
- la Fimbristylis cymosa
- la Euphorbia prostrata
- pl Cucurbita pepo
- pl Plumeria rubra
- pl Gomphrena globosa
- ec Lycopodium nodiflorum
- c Miconda citrifolia
- ec Portulaca oleracea
- pl Zephyranthes rosea
- ec Pilea microphylla

- pl *Asclepias curassavica*
 c *Flemingia strobilifera*
 lc *Paspalum conjugatum*
 a *Nephrolepis hirsutula*
 a *Polypodium scolopendria*
 pl *Veitmania sanctum*
~~*Asclepias curassavica*~~
 cla *Scaevola sericea*
 la *Tournefortia argentea*
 pl *Taqates*
 pl *Hibiscus mutabilis*
 l *Clerodendrum inerme*
 pl *Hymenocallis littoralis*
 pl *Cinnamomum asiaticum*
 pl *Polyscias triochloata* (chloata)
 lc *Allophylus timorensis*
 lc *Calophyllum inophyllum*
 cla *Asplenium nidus*
 lc *Hemodia tuba*
 lc *Guettarda speciosa*
 n *Persea samboensis*
 n *Boerhavia tetrandra*
 lc *Intsia bijuga*
 n *Vittaria elongata*
 lc *Nephrolepis acutifolia*
 la *Elichraia geniculata*
 c *Alocasia macrorrhiza*
 lc *Hibiscus tiliaceus*
 la *Boerhaavia gongilodes*
 n *Ophiorhiza pendulifera*
 n *Pteris tripartita*
 pl *Cordia alliodora*
 pl *Acalypha wilkesiana*
 pl *Prenderanthemum cantharidis*
 var. *atropurpureum*

Pipturus trees that are
 chlorotic have very few red mites.
 2-3 *Zoysia tenuifolia* supposed
 to have been brought from
 Java, to which it was
 introduced by the Japans.

Monarch butterflies and
 larvae abundant around
 mulweed patch in village.

The north part of islet
 brushed out for planting
 in December.

Whimbrel on day pass reef.
 White-capped noddy.

Mangrove swamp near
 north end almost clear.
 Scattered clumps of
 very slender *Bruguiera*
 to 5 m. tall. Many
 seedlings. Has been
 cleared somewhat to plant
 coconuts. Many volunteer
 coconut seedlings. Many
 fallen and standing trunks
 covered by epiphytic ferns.
 This cleared swamp is
 adjacent to an open-water
 pond, strongly brackish,
 great masses of blue-green
 algae of a rusty yellow-green
 color. Apparently rather
 little tidal fluctuation.
 A few big *Bruguiera* around

frond, some saplings,
mainly coconut seedlings
Eleocharis in mud
back from edge. Coconut
seedlings somewhat
a very chaotic.

Much trash in wood.
~~Several~~ fish about 20-25 cm
long, mud color with
black fins yellow tail,
mullet shape.

Several white terns,
one or two tattlers.
White capped noddies.

~~and~~ golden plover.

Red cellula, large.

one or two common noddies?

Big greenish Acrochorda gomphe
Large area of this
swamp, mostly semi-
open. Around this,
especially southward,
is a tall scrub or low
forest of Pipturus argenteus,
slender, 5-6 cm dbh maximum,
to 4-5 m tall. Has been cleared
beneath. Varies from
soil of some depth to
boulder field. Ferns
abundant.

In interior are considerable
open areas with scattered
breadfruit trees, covered by a
blanket of Wedelia.

- o Premna obtusifolia
- pl Tolypsis scutellaria
- pl Citrus aurantifolia
- l Euphorbia chamissonii
- l Cyperus javanicus
- lo Cyperus compressus
- o Physalis angulata
- lo Ipomoea littoralis
- ea Cyperus odoratus
- pl l Cyrtosperma chamissonii
- pl l Colocasia esculenta
- l Jussiaea suffruticosa
- c Canavalia microcarpa

Other areas have grown
up to the low forest type
dominated by Pipturus.
In places this is cleared
out locally. Some areas
the average height is
6-8 m.

The breadfruit trees injured
by the typhoon either died or
have recovered remarkably
well. Branches as large
as 10 cm diam. have grown
since the storm, crowns
well filled out, symmetric.
Leaves dark green.
Some infestation of cottony

cushion scale. Some trees have died of a peculiar disease. (Kruskin describes it in report to Hi Com.)

Taro pit - generally dominated by *Thelypteris goepeloides*, locally by *Cyperus odoratus* or *Echinochloa*, locally invaded by *landanum*, *Vigna*, etc. *Jussiaea* common.

Very little taro cultivation - two rows and scattered plants of *Cyrtosperma*, scattered small patches of stunted *Colocasia*.

In other pit *Thelypteris* even more generally dominant, almost exclusive over most of the area. Small patches of *Eleocharis*, *Sporocytus tuba*, open water with blue-green algae. Some invasion by *Pandanus*. Occasional *Jussiaea*.

No *Echinochloa*. Only a very small part of this pit occupied by *Cyrtosperma*, no or very little *Colocasia*.

Of total area of two pits perhaps not more

than 1% occupied by either taro. Only a few sq. m. cultivated - *Cyrtosperma*.

Aedes common.

Wedelia blankets all openings in this vicinity, incl. margins of taro pits.

Paspalum conjugatum is very locally common along trails in semi-shade, nowhere really abundant.

Mangrove swamp between taro pits and village has all larger *Brucea* trees dead and broken off part way up. Ascribed to typhoon.

Many well grown seedling coconuts generally. *Pandanus* trunks mostly rather well rotted, coconut less so.

Along seaward side a well-developed *Scaevola* fringe 2-3 m. tall.

Epiphytes common to abundant on older trees in interior.

Oct. 29 - Lunglap Islet,

- 41405 *Thelypteris goggholodua*
dominant in semi-abandoned
taro pit.

- 16 ✓ 06 *Jussiaea suffruticosa*
common in ^{semi-abandoned} taro pit

- 3 07 *Echinochloa crusgalli* (L.) Beauv. ^{semi-late}
locally dominant in
semi-abandoned taro pit

- 12 ✓ 08 *Zoysia tenuifolia*
locally planted in home
yards of coral gravel

- 2 ✓ 09 *Ophiosphragis pendulum* L.
rare, epiphytic in
wet forest

Oct. 20 - Kirajon I.

- 1 ✓ 10 *Allophragma timorensis* Bl.
common in dense scrub on grass
beach ridge.

- 1 ✓ 11 *Eutadea purpurea* DC.
one plant on beach ridge.

Oct. 26 - Imroj I.

- 4 ✓ 12 *Vigna marina* (Burm.) Merr. ^{W.L.}
abundant on new gravel

Oct. 23 - Ribon I.

- 7 13 *Vittaria incurvata* Cav.
rare, epiphytic on tree trunks

erect from rhizome buried
slightly in mud.herb 1 m. tall, branched;
petals 4, yellow, caducous
erect.

forming a close dense mat.

sterile

large shrub 4 m. tall,
seedlings abundant beneath
it.
seedling

unifoliate

prostrate, sterile; note
unifoliate leaves, these
seem to be characteristic
of rapid-growing shrubs
at least in this area.

Oct. 26 - Trip, Jabuon to Imruj
Crested tern flying over
lagoon.

Oct. 26 - Imruj Islet

Plants seen

- pl
a
pl
c
pl a
pl
pl
a
pl
o
pl a
pl a
pl a
pl
a
pl a
pl a
pl a
pl
a
pl a
pl a
pl a
bc
o
c
pl
la
c
c
la
c
la
c
- Gynophrena globosa*
Polypodium scolopendria
Plumiera acutifolia
Carica papaya
Musa sapientum
Zinnia elegans
Mirabilis jalapa
Crocos nucifera
Galearias curassavica
Lissonia grandis
Citrus aurantifolia
Hibiscus hybrid (very clustered)
Cordia terminalis
Cucurbita sp.
Centella asiatica
Ipomoea batatas
Cucumis sativa
Barringtonia asiatica
Physalis angulata
Morinda citrifolia
~~*Xanthosoma sagittifolia*~~
Tournefortia argentea
Eleusine indica
Artocarpus altilis
Digitaria pruriens v. *microbachne*
Pandanus tectorius
Pipturus argenteus
Alocasia macrorrhiza

Bananas very luxuriant
around house.

Centella is remarkably
abundant on gravel sheets
in full sun.

Young *Tournefortia* 2-3 m tall.
Sweet potato patches
very common.

Phosphate bed covered
by an open scrub of
Pipturus with some
areas closed, some *Clerodendron*
much *Lepturus* and *Digitaria*
filling interstices.
Flemingia and *Polypodium*
prominent on rocks,
here and there *Wedelia*
mats, a little *Nephrolepis*.

Toward the pass this
becomes ^{locally} closed, 3 m tall,
and composed of *Scaevola*,
Tournefortia, *Guetarda*,
Barringtonia (from old stubs)
Thaarea and *Pipturus* in
openings.

Two former mangrove
depressions at north end -
one is completely bare, the
other has many *Prugniers*
seedlings and a few larger
but still small plants.

1 grayish mottled reef heron

Plants seen (old.)

- pl *Cinnamomum asiaticum*
 pl *Hymenocallis littoralis*
 pl *Polyscias scolotheca*
 o lc *Cenchrus echinatus*
 la *Lepturus repens*
 pl *Angelonia ^{angustifolia} angustifolia?*
 pl *Cathara thus rosea*
 pl *Zephyranthes rosea*
 la *Fimbristylis ^{cymosa} cymosa?*
 a *Vigna marina*
 lc *Hedyotis biflora*
 o *Calophyllum inophyllum*
 la *Leacvola sericea*
 o *Eragrostis amabilis*
 o la *Pennisetum ruderale*
 lc *Thuarea involuta*
 a *Wedelia biflora*
 c la *Triumfetta procumbens*
 lc *Clerodendrum inerme*
 lc *Nephrolepis bisentata*
 c *Quettarda speciosa*
 a *Peperomia ponapensis*
 a *Euphorbia charmissonis*
 pl *Cordia allamanda*
 c *Allophylus timorensis*
 la *Bruguiera gymnorhiza*
 lc *Kalanchoe pinnata*
 a *Caesalpinia*
 lc *Ipomoea littoralis*
 lc *Synedrella nodiflora*
 a *Vernonia cinerea*
 pl *Rhoeo spathacea*
 pl *Canna hybrid*

Broad gravel ridge
 running parallel to depression
 is still partly bare
 but much colonized by
 fair sized *Triumfetta*,
Leacvola, large ^{low} mats
 of *Lepturus*, a few of
Triumfetta, locally
 abundant seedling
 of *Euphorbia charmissonis*
 meeting clumps of *Allophylus*
 some *Thuarea*, seedlings
 of *Triumfetta* + *Pennisetum* ^{seedling}

(top 72)
 In central part,
 back from lagoon
 beach, is an open
 forest, mostly *Pisonia*
 8-10 m. tall, some *Antrocarpus*,
 some *Allophylus*, some
Pipturus. A great deal
 of dead brush, obviously
 chopped *Pipturus* which
 must have formed
 a dense scrub 3-5 m.
 tall. Sprouts from
 stubs of this, and from
 other species also, esp.
~~some~~ *Pisonia* and *Allophylus*,
 abundant, 1-1.5 m. tall.
 This is on small gravel.
 Seedling breadfruit to
 5 m tall common.

Oct. 26 - Kinajon I.
Plants seen

- l Portulaca oleracea
pl Oenothera sanctum
la Phyllanthus amarus
lc Cyperus javanicus
o Alcocasia macrochiza
la Euphorbia hirta
c Carica papaya
c Artocarpus altilis
la Lepturus repens
a Cocos nucifera
a Polypodium scolopendria
r Pteris tripartita
la Nephrolepis bisectula
a Pipturus argenteus
a Vigna maritima
pl Xanthoxylum sagittifolium
a Wedelia biflora
pl Bougainvillea
c Pandanus tectorius
pl Musa sapientum
lo Cordia subcordata
lc Morinda citrifolia
c Quettarda speciosa
pl Cucurbita
la Cassytha filiformis
o Physalis angulata
lc Trumpetta procumbens
a Fimbristylis cymosa
pl Pseudanthemum carthagenum
var atropurpureum
etc Jacca leontopetaloides
r Pilea microphylla

Planted area back of house is dominated by Nephrolepis locally Vigna, Polypodium, becoming overgrown by Wedelia. Some spots still covered by Fimbristylis and/or Lepturus. Farther back toward sea Polypodium and Lepturus are codominant, a little Wedelia. (Photos)

✓ Fairy tern. Common noddy. Many volunteer coconuts. Strip along beach is mixed scrub - Scaevola, Tournefortia, Quettarda, a little Allophylax, tangled in places with Canavalia and Wedelia occasional Cordia. This strip varies, up to 40 m. wide, 2-4 m. high. Two home sites, one at landing on lagoon shore one opposite on seaward side.

Gravel sheet opposite rounded indentation on seaward side - blackened by algae. Colonized by seedlings of Quettarda, Scaevola, Lepturus.

Plants seen (ctd.)

- lc *Tournefortia argentea*
- e *Allophylus timorensis*
- l *Canavalia microcarpa*
- la *Thuarea involuta*
- lc *Calophyllum inophyllum*
- r *Ochroma oppositifolia*
- e *Asplenium nidus*.
- o *Pisonia grandis*
- la *Sporaea tuba*
- r *Nephrrolepis acutifolia?*
- l *Hibiscus tiliaceus*
- l *Kalanchoe pinnata*
- l *Barringtonia asiatica*
- r *Terminalia samoensis*
- r *Cleome indica*

Morenda, Pandanus,
Triumfetta, Wedelia.

Morenda very abundant
in inner part. Some
Polypodium.

Ragged but well-recovered
Calophyllum trees scattered
along top of gravel beach
here.

Where shore bulges
out, south of bay, there
has been some clearing
out of brush, resulting
in a scattered stand of
small ragged trees,
tangles dead brush,
coconut seedlings.

Intrisia seedling on
gravel ridge.

South on curve most
exposed to sea, is a dense
~~or~~ tangled scrub or
scrub forest of *Allophylus*,
Guettarda, *Calophyllum*,
Boerhaavia an *Alchornea*
sapling a little *Tournefortia*,
but dominantly *Allophylus*.
This on beach ridge of gravel.
Asplenium nidus common
here.

Seedlings of *Allophylus*
abundant, *Guettarda* less
-11-

This scrub is apparently left as a wind break on the beach ridge of medium poorly sorted gravel, as it is only a few m. wide. Inside the coconut plantation has been cleaned of brush, but is regrowing it.

Large mangrove depression has older trees, to 8 m. tall rather scattered, ~~but~~ middle layers about 1-2 m. also sparse, lower layers, 0.5 m. mostly dense.

Where brush is cleared in interior *Wedelia* is growing from stumps, also millions of *Wedelia* seedlings.

Scrub belt very dense on outer beach ridge changes to *Wedelia* with bushes of *Tournefortia*, *Pipturus*, etc. 2 m. high as coast swings round toward south passage beach.

White reef heron. Noddies.

Coconut plantation recently cleared in from low. Many young trees, but probably dating from before typhoon. This low belt of scrub continues around on beach ridge on passage beach. Here platform of breccia is very well exposed. (photos below).

Inside this beach ridge is a swamp that was almost completely cleared of *Bruguiera* by the typhoon - is still open except for a few small *Bruguiera* saplings. Bottom is rather soft mangrove peat with a layer of fine coral gravel on top of it. Enormous mass of rotting debris.

Inside this is a concentric ridge with sparse coconut trees, then another *Bruguiera* swamp, this one not so damaged in part, part completely ^{clear}.

In outer swamp, about half way from edge to pool in center the peat is 9" deep, with about an inch of mixed sand, gravel and peat on top. Below is gravel.

(Samples # 21, 22)

Ridge bet. swamps
cleared and staked but
not yet planted to coconuts.

Lagoon end of inner
swamp open, with only
a few living small
Bruguiera, old trunks
lying all in one direction,
to interior of islet.

(photos of this and
cleared ridge)

Lagoonward end of
outer swamp half filled
by a thick gravel sheet
that came in from the
seaward quarter of
passage beach, ending
abruptly, with an
irregular steep slope
about 1.5 m. high, top
very flat. Bottom of
swamp here of sharp
irregular boulders.

This gravel sheet
lines the coast, extending
well around on passage
beach proper, narrowing
to a typical boulder ridge.
Its top is in part covered
by Wedelia, in part
by widely spaced
Tournefortia bushes
about 2-2.5 m. high.
Some Scaevola, part open

and blackened.

Several white tern
and white-capped noddies.

Inside the second
mangrove swamp is a
thicket of *Sonneratia*
tiliacea, then a third
more extensive *Bruguiera*
swamp, the taller trees
dead, but a thicket of
smaller ones, 3-4 m. tall.

Much debris in margin,
esp. old *Bruguiera* poles,
all lying one direction
more or less southwestward,
not quite same as ^{some of} coconuts
lying in swamp which
are more south. Much
rough cobble size
coral gravel + boulders
swept into swamp
here from passage
beach as edge of thinner
gravel sheet, overlying
mangrove peat and
sinking into it when
stepped on. In swamp
on a few older trees are
a few epiphytic *Asplenium*
nidus, *Polypodium scolopendria*, and one *Nepenthes*
acutifolia (?) (too young or
at least small, to be sure
it is this sp. but probably).

Around edges of swamp on sloping steep coral gravel, seedling *Bruguiera* about 3-4 dm. tall abundant to high tide marks, but do not look especially healthy. Around edges, above h.t. are patches of *Spartina*. Then, extending to lagoon, a blanket of *Wedelia* at least 1-m. thick, partly cleared inland, with scattered trees, incl. tall *Pisonia*, lower *Allophylus*, *Guettarda*, *Pipturus*, *Pandanus*, and *Coco*.

^{p 75} Land staked for planting but not planted.

East of last swamp is another pond, only a few m. across, with 2 or 3 *Bruguiera* not more than 1-1.5 m tall. *Hibiscus* tiliaceous tangle on one end.

(insert here part from p 69, put there by mistake)

Boulder + small grass ridge lying on lobe

At Beccia platform on lagoon side, has grove of small *Barringtonia* (photo below). *Terminalia samsonensis* occasional along lagoon gravel ridge.

Weathered spathe of spicate coconut on lagoon beach.

Aedes occasional.

Lagoon shore and back of it toward north pass much more sandy than elsewhere.

Oct. 26 - Monday?
 north, southward from
 landing.

Bananas very dense
 and in good shape,
 few coconuts.

Then sweet potato
 area - boulders and
 coarse gravel piled
 up to clear sweet potato
 patches.

Then open gravel
 sheet, not yet cleaned
 off for planting.

Tall Scaevola, Pipturus,
 Tournefortia, Wedelia
 scrub.

Then high ground
 planted to coconut
 seedlings, otherwise
 grassy toward lagoon
 to m. scrub toward sea.

This tapers off toward
 end of islet into Wedelia
 mat with scattered
 Pipturus, Tournefortia,
 Scaevola

This changes to
 an open scrub of same
 species, tangled with
 Wedelia

1 whorled ones 1 taller

Plants seen (ctd.)

Pseuderanthemum cantherosii
 var. *atropurpureum*

Pteris tripartita

Cordia subcordata

Cassytha filiformis

Ochroma oppositifolia

3 fairy terns on outer
 reef.

On outer edge the
 scrub is very ~~open~~
 and about 1 m. high.
 Some *Triumfetta* and some
Wedelia on ground between
Scaevola, *Tournefortia*
 and (a) *Quercus* bushes
 some on ground. Some
Pandanus seedlings.

Some patches of *Lepturus*.
 This character is
 maintained for some
 distance north along
 seaward beach, but
 locally the bushes are
 2 m. tall

Hernandia, *Allophylus*,
Pandanus seedling, rare on
 fresh gravel ridge at
 top of beach. Also *Calophyllum*,
Scaevola, *Vigna*.

Some *Vigna* appears
as well as northward.

This ~~part~~ reasonably
fresh beach-rock at
about mid-tide,
hardened and the surface
abraded, but appears
quite fresh. (sample # 23).
Sample # 24 is sand from
the upper part of beach-
a rather thin deposit.
for examination as to
nature of material. Seaward
beach near south end.

north from landing

Bananas around village.
then sweet potato patches,
then an open scrub
of *Pipturus*, *Pandanus*,
much *Wedelia*, some
Digitaria patches, *Vigna*
some *Tournefortia* and
Scaevola.

This gives way abruptly
to a grassland of *Thuesia*,
Lepturus and *Digitaria*
with scattered coconuts,
all sizes, and tall bushes
of *Premna*, *Tournefortia*,
Pipturus, *Allophylus*,
Pisonia, *Terminalia samoensis*,
clump *Musa* of *Alouasia*
Seaward is a tangle

of *Wedelia* and shrubs,
tapering northward to
a beach ridge with
Tournefortia.

The grass becomes
more and more dominant
northward, with few
shrubs except young
coconut and an occasional
Morinda. In places
Polyposidium almost
replaces grass. *Fimbristylis*
and *Plumiza* common.

Along lagoon beach
occasional battered
and distorted, (but sprouting
abundantly) *Cordia* trees.

Small colony of *Lechona*
on beach ridge at end
of inlet, with *Scaevola*,
Tournefortia,
about 4 m. tall, getting
lower ~~se~~ and more open
seaward, with seedlings
and sprouts of *Barringtonia*
in grassy opening.

Two almost white
reef herons in mangrove
depression.

Wandering tattlers on both
beaches and in mangrove
depression.

Oct. 27 - Lijerua Islet.

Species observed in 1958:	1960
<i>Asplenium nidus</i>	✓
<i>Lepturus repens</i>	
<i>Cocos nucifera</i>	✓
<i>Pisonia grandis</i>	✓
<i>Canavalia microcarpa</i>	✓
<i>Intsia bijuga</i>	
<i>Vigna marina</i> (2)	
<i>Galaplythum inophyllum</i> (2)	
<i>Pemphis acidula</i>	✓
(27) <i>Terminalia samoensis</i>	✓
<i>Barringtonia asiatica</i> (2)	
<i>Cordia subcordata</i>	✓
<i>Tournefortia argentea</i>	✓
<i>Quettarda speciosa</i>	
<i>Scaevola</i>	✓
<i>Pandanus</i> (2)	✓

Birds

	Frigate bird	100-200?
nesting	White capped noddy	many
one egg	white tern	faintly common
	Crown booby	a number
	Bristle-thighed curlew	2 or 3
	turnstone	medium flock
	tattlers	one or several
	plovers?	
	Reef heron	1 speckled

Some of boobies were white beneath, only seen flying. could not see other features.

Islet presents a solidly vegetated appearance low forest with a few coconuts and several other taller trees as emergents. Pemphis patch still presents a weatherbeaten appearance but is generally leafy.

Generally a mass of *Pisonia* forest with complete canopy at perhaps 8 m. mainly of sprouts from fallen trees. Some standing trees with tops broken and masses of sprouts, others standing out as emergents.

Around margins *Terminalia* and *Tournefortia* are common, forming a zone. One clump of *Scaevola* 2.5 m. tall in this, a small tangle of *Canavalia* here, also, with the *Scaevola*, fruiting. A few standing coconut seedlings and young trees. One fallen *Cordia* with abundant sprouts on beach.

Asplenium common on fallen trunks and ground many very old dead bases. locally forming a ground cover.

Humus beds under
fringing well developed
to 5-15 cm thick. Very locally
up to 1 cm of poorly consolidated
phosphate rock.

Tournefortia seedling
common on n. sand flat.

1 green turtle nest 130 eggs.

Logs and branches are much
more rotted than on other
islets, possibly from Ni-
guans.

Two long sand horns
extending north and
south from lagoon corners.
Between these active beach
erosion is taking place,
undercutting the shrubs
on shore.

Oct 27 second islet north of
Mejatto in bare sand
with about 2 coconut trees
and a few bushes.

From lagoon both ends
of Mejatto appear densely
vegetated, quite a few
coconuts, thick undergrowth.
Between, very sparse
coconuts, low scrub
undergrowth.

Oct. 27, Mejatto, ^{from} north end,
to south.

Plants seen

- 20 *Premna obtusifolia*
- 2 *Anticarspus altalis*
- 20 *Lepturus repens*
- pl *Cucurbita*
- pl *Carica papaya*
- pl *Pomoea batatas*
- 20 *Pandanus tectorius*
- 20 *Digitaria pruriens* - *umbellata*
- 1 *Portulaca oleracea*
- pl *Mirabilis jalapa*
- pl *Crinum asiaticum*
- pl *Alocasia macrorrhiza*
- pl *Clerodendrum inerme*
- 20 *Pleroma ruderalis*
- pl *Plumeria rubra*
- pl *Musa*
- 2 *Wedelia biflora*
- 20 *Fimbristylis cymosa*
- 1 *Polypodium scolopendria*

- o *Morinda citrifolia*
 n *Nephrolepis hirsutula*
 la *Thunaea involuta*
 n *Tacca leontopetaloides*
 la *Vigna marina*
 o *Eleusine indica*
 a *Scaevola sericea*
 a *Pipturus argenteus*
 c *Zuetharda speciosa*
 n *Ochrosia oppositifolia* (S)
 lc *Allophylus timorensis*
 n *Intsia bijuga* (S)
 lc *Sporobolus tuba*
 r la *Triumfetta procumbens*
 o *Terminalia samensis*
 lc *Barringtonia asiatica*
 lc *Pisonia grandis*
 lc *Physalis angulata*
 a *Tournefortia argentea*
 lc *Cordia subcordata*
 lc *Cassytha filiformis*
 lc *Canavalia microcarpa*
 n *Centella asiatica*
 n *Pseudorantherium* var. car.
 2o *Calophyllum inophyllum* var. car. "luquej"
 n *Pemphis acidula*
 n *Hernandia sonora* (S)
 n *Borhavia tetrandra*
 n *Euphorbia cheimissonis*
 179 la *Bruguiera gymnorhiza*
 n *Nephrolepis acutifolia*

north end of Mejatto large clearing, grassy with *Digitaria* and some *Lepturus* and *Fimbristylis*, scattered coconuts, many bread-fruit saplings & seedlings, up to 2.5 m. tall.

This extends perhaps half way to the sea. Then a dense tangled thicket of *Pipturus*, *Scaevola*, etc. about 3-4 m. tall, with *Wedelia*. This with fallen coconut trunks this extends to near the outer beach, where it thins and becomes lower, becoming open on beach ridge. Here the beach ridge is of cobbles & boulders, perhaps 2.5 m. high.

A few *Terminalia* seedlings on bare beach ridge. *Sporobolus tuba* running out onto it. Extensive rubble tracts on reef flat.

Flocks of 6 white terns fishing just outside outer reef. 2 white reef herons, one gray & white one.

Southward from here on lagoon shore the old clearing becomes more and more overgrown with *Vigna* and *Wedelia*, and the space between the ~~the~~ scrub belt and lagoon beach narrows.

Then, where islet abruptly narrows down, wide areas are covered by a blanket of *Vigna*, covering a sandy gravel flat. This extends along the lagoon coast a bit, changing inland to an open scrub of *Pipturus*, *Tournefortia*, etc. tangled with *Vigna* and *Wedelia*.

Along lagoon beach and reef flat there are wide, rather thin rubble sheets with a lot of old tree trunks and rubbish.

South of the *Vigna* area is a flat of gravel with scattered *Tournefortia* bushes 1-1.5 m. tall, then *Lepturus*, thin *Vigna*, thin *Wedelia*, presenting a rather arid aspect. Appears

to be an area scoured by typhoon. Inland a coarse gravel sheet with lobes margin, flat, much of it open, scattered *Tournefortia* and *Scaevola* about 2 m. tall, ground either bare blackened gravel or thin *Wedelia* mat, some *Triumfetta*.

Seaward *Tournefortia* become lower *Wedelia* denser and thicker, forming a closed vegetation. Become open again and irregular grassy, with *Triumfetta*, open gravel, *Vigna* and scattered low *Tournefortia* and *Scaevola* in outer 50-75 m. A low bare gravel ridge outside this. Steep gravel beach about 2 m high. Then a rubble tract running perhaps half the length of the island. photos

This rubble tract, uncovered at low tide, has several sandy strips across it, perpendicular to beach.

On the gravel sheet back of the beach ridge about half way to where the open scrub starts is

a line of debris - old husks, sticks, coconuts, etc. on which a dense low stragg of vegetation about 1-3 m. wide has developed - mostly *Scaevola* up to 0.5 m. tall, numerous young coconuts, little taller, *Canavalia*, *Triumfetta*, *Cassytha*. Curiously no *Tournefortia*, though it is abundant, dominant, in the ^{open} scrub inland and as numerous as *Scaevola* (seedling) in the gravel sheet seaward to the gravel ridge.

Standing coconuts are very sparse in this section of the islet.

Gravel sheet rather thin here, old coconut root masses occasionally protrude through, covered with *Lepturus* and *Fimbristylis*.

A little farther south, at perhaps the narrowest part of the islet the outer portion - gravel sheet, plunge holes, scoured areas, etc. has only a sparse mat of *Triumfetta* and *Lepturus*, some *Vigna*, *Sporobolus*, etc.

Wedelia, etc. with scattered very small *Tournefortia* about 3-5 dm. tall. *Fimbristylis* patches very locally.

This thickens lagoonward to a scattered scrub of *Tournefortia* in a blanket of *Wedelia*. South of this an area of tall dense scrub of *Cyrturus*, *Tournefortia*, *Scaevola*, *Wedelia*, etc. Then a scoured open area of *Thuarea*, *Lepturus*, *Triumfetta*, *Polypodium*, etc. with scattered shrubs. Southward, again, this thickens to an open scrub and *Wedelia* blanket.

Outside ^{of this} the gravel sheet in from the gravel ridge, along all of this, is covered by a sparse scrub of *Tournefortia*, *Scaevola*, some young coconuts, etc. with patches of *Lepturus*, *Triumfetta*, etc.

Gravel ridge has white. Southward this ^{open} ~~area~~ scrub strip narrows considerably, the low scrub, with *Wedelia* or *Lepturus* extends ^{almost} the width of the islet, dominated

mostly by *Tournefortia*, locally *Pipturus* or *Scaevola*. Some open gravel spaces. *Vigna* toward lagoon. Southward much open *Thuarea* and *Lepturus*, or *Vigna*, with scattered *Pipturus* bushes, 2-3 m tall, rounded, and *Tournefortia*, 1-2 m tall. Inland a large bare gravel sheet of slabby boulders, rather blackened, parts of this with old coconut trunks lying on ground and scattered shrubs. Seaward a narrow strip of thicker lower scrub tangled with *Wedelia*, then the typical sparse *Scaevola* *Tournefortia* about 3-4 dm tall of the seaward gravel sheet and ridge, some *Guettarda* and scattered stunted coconut seedlings.

This strip across is about $\frac{1}{3}$ the length of the islet, north to south and has almost no standing coconuts, but

some stubs. Just south of it taller and denser scrub, considerable *Vigna* on seaward side, the open low scrub area very narrow.

Then a badly scoured area across the islet, much open gravel, scattered bushes, some grass, *Thuarea*, *Pipturus*, some *Wedelia* & *Triumfetta*, scattered *Tournefortia*, *Pipturus*, *Scaevola* shrubs 1-2 m tall. This extends some distance scrub and few gravel in varying proportions. Thinning out and becoming more desolate in appearance southward, with scattered shrubs mostly about 1 m tall, of *Tournefortia*, patches of *Lepturus*, *Wedelia*, *Vigna*, *Thuarea*, etc., but mostly bare gravel with coconut roots crumbled lagoonward, old bases, fallen trunks, etc. Many plunge holes. This area shows only the beginnings of recovery. Bounded southward by a

depression running across the islet with practically no vegetation at all.

except along north side a scattering of seedlings of *Scaevola* and *Tournefortia*, a very few around a couple of old coconut bases in the middle. Across the seaward side is a gravel ridge. An lagoon reef flat is a pond surrounded by a circular gravel bar.

pt
1 crested terns on lagoon beach.
2 blue herons lagoon beach.
1 white on seaward beach.
2 tattlers on seaward reef flat.

Several enormous dead *Calophyllum* on lagoon beach - uprooted.

Oct. 27 - I mung
along seaward beach
from north end -

Low open scrub of *Scaevola*,
Tournefortia, *Peperomia* with
mats of *Trunfetta*,
Peperomia, *Thunbergia*, some
bare gravel.

Varying greatly ~~to~~
from place to place in
both density and height.

4 turnstones and a plover.

Oct. 27. Ribbon I.

Seedlings on gravel bar
Scaevola, *Pandanus*,
Calophyllum, *Quettarda*,
Marringtonia, *Morinda*,
Cordia, *Vigna*, *Tournefortia*

Plants seen

- l *Cassytha filiformis*
 a *Asplenium nidus*
 a *Wedelia biflora*
 a *Pisonia grandis*
 c *Polypodium scolopendria*
 cc *Cordia subcordata*
 c *Morinda citrifolia*
 n *Vittaria elongata*
 cc *Canavalia microcarpa*
 cc *Tournefortia argentea*
 n *Nephrolepis acutifolia?*
 l *Flemingia strobilifera*
 o *Scaevola sericea*
 la *Vigna marina*
 c *Quettarda speciosa*
 2 cc *Calophyllum inophyllum*
 4 cc *Pandanus tectorius*
 2 cc *Marringtonia asiatica*
 cc *Cocos nucifera*
 la *Pemphis acidula*

1 bristle-thighed curlew,

eggs, also

Flock of 15-20 black-naped terns around end of gravel bar, one fully grown young, 2 common noddies, and quite a few white terns. One almost grown young in tree. *Aedes triseriatus*.

Bottom of enclosed pond now about half covered by low *Pemphis* scrub perhaps 1.5 m. tall. some new seedlings or sprouts from old roots. A little *Lepturus*, some *Canavalia*. Bar surrounding it largely covered by scrub of *Scaevola* with a little *Tournefortia*, some *Canavalia*. This varies from 1-2.5 m tall. A little *Terminalia*, *Vigna* around edge.

Plants seen (ctd)

- cc *Lepturus repens*
 c *Terminalia samoensis*
 n *Pemphis argentea*

and south
 The west ~~ends~~ present
 a solid mass of vegeta-
 tion, largely *Pisonia*,
 up to 10 m tall, maximum,
 mostly 5-6 m. *Cordia*,
Scaevola, *Morinda*
 occasional *Tournefortia*,
Guettarda. Tangled
 with *Wedelia* and occasional
Canavalia.

Interior ~~is~~ dominated
 by *Guettarda* and *Pisonia*
 sprouts about 4 m tall,
 with an undergrowth
 of *Asplenium nidus*
 2 m tall on large
 root masses. Some
 of these old root masses
 dead, but covered by
 countless young *Asplenium*
 plants. There also numerous
 or old coconut bushes,
Loq, etc. *Asplenium*
 also epiphytic in lower
 parts of trees, as is *Polypodium*

A few Coconut trees in
 north part of islet.

This part generally
 is a low irregular scrub
 mostly *Pisonia* sprout
Scaevola and *Guettarda*
 tangled with *Wedelia*.

Here a small colony of
Flemingia on gravel
 under a *Pisonia* tree.
Wedelia on base of bar.

On base of west bar
 a scrub of *Scaevola*
 and *Guettarda*. Distally
 from this, a mat of
Casuarina, with
 many seedlings of *Pandanus*
 and *Calophyllum*, *Barringtonia*.
 Distally from this a mat
 of *Vigna*.

^{on islet}
 Here the largest *Pandanus*
 is a single stem about 1 m or less
 tall, others very small seedlings.
Calophyllum, *Barringtonia*
 only seen as seedlings.

This islet has become
 very luxuriantly overgrown
 up to about 5-6 m, with
 coconuts and a few *Pisonia*
 trees as emergents.

28. 28 - Mejatto, north point
and passage beach.

Well developed *Scaevola*
fringe, up to 3 m. high,
along passage beach.

Back of this a sheet of
coarse gravel with
a few scattered bushes
of *Scaevola* & *Tournefortia*
an old *Tournefortia*,
scattered mats of
Cassytha, *Wedelia*,
& little *Triumfetta*,
very sparse tufted
Lepturus, this especially
but not exclusively, on
sandy patches. *Cassytha*
on *Wedelia* and *Lepturus*,
scattered seedlings
of *Scaevola*, *Tournefortia*.

This was formerly a
sandy area where coconuts
were always yellow, as
the boys who live here.

Toward sea the
Wedelia and *Cassytha*
become abundant
and the ground cover
complete. This coincides
with end of gravel
sheet and greater
sandy area of ground.
Lepturus and *Fimbristylis*
also more abundant, and *Plum*.

Toward seaward
corner the *Scaevola* fringe
is interrupted by a wide
sheet of coarse gravel,
partly low, partly
covered by an irregular
Scaevola scrub
fading inward to a
mixed open scrub
with *Pipturus* and *Wedelia*.
Passage beach lined generally
by a low narrow ridge of
pebble gravel.

- 1 white heron
2 turnstone
1 white tern

Middle of Mejatto where
I left off yesterday
south of big scum
an area is being colonized
by *Scaevola* and some
Tournefortia, a few stunted
yellow coconuts, none of them
over 0.5 m. mostly less,
sparse. Toward lagoon
on remnants of soil
is irregular scrub &
grass. A few standing
coconut trees and many
stubs.

Then for a stretch there

is an irregular, very low scrub, quite sparse. remnants of soil filled with coconut roots. small thin mats of *Triumfetta*, patches of *Lepturus*, seedlings of *Landanum* + *Barringtonia* but main components *Scaevola* + *Tournefortia*. A few seedlings of *Quettarda*, *Hibiscus tiliaceus*, *Mouroua*, *Wedelia*, *Pemphis*, *Calophyllum*.

Then another deep scour.

The entire area between them, a deep scour channel has been scoured off except for remnants of soil held by coconut roots. (b-w photos)

The second scour also has a gravel ridge on seaward side, not so complete. No vegetation except on several areas of firm soil not entirely scoured off. Here a sparse *Scaevola*, ~~the~~ *Tournefortia* scrub 0.5 m. tall.

Southward after is broken up area of soil remnants and small

scour channels, with ~~at~~ very irregular mixed scrub and grass patches, a taller mixed scrub of *Tournefortia*, *Pipturus*, *Scaevola*, etc. (b-w photos)

This very open, with patches of grass (*Lepturus*) and of open gravel sheet. Bushes rounded. 1-3 m. tall.

(*Pipturus* = *arumua*, acc. Ellis)
(used for cordage.)

Southward, shrubs more open. *Signa* mats on lagoon side, *Thruarea* and *Wedelia* seaward.

Small mangrove swamp, sand bottom, with irregular but pure stand of *Bruguiera* 1-1.5 m. tall. near lagoon, separated from lagoon by sand ridge, encroached upon from seaward by gravel ridge. No larger *Bruguiera* trees, but a few dead skeletons lying lagoonward, pointing toward lagoon.

Gravel sheet to seaward, very sparse scrub, mostly *Pipturus*, with grass mats and ~~the~~ gravel. *Fimbristylis* (photo)

1 flower
1 stem

seedlings locally abundant.

Lepturus + *Thuarea* more abundant southern on gravel and on sand ridges along lagoon.

Local patches of *Euphorbia chamoissonii*, no obvious reason. Some *Triumfetta*

then tall open scrub of *Pipturus* + *Tournefortia* with grass, *Triumfetta*, *Wedelia* + open gravel patches.

Crested tern on lagoon side. harsh squawking etc.

Tall scrub becomes lower, 1-1.5 m. tall, with solid grass, *Lepturus* + *Thuarea*, ground layer occasional mats of *Wedelia*.

bearded leaves, more open, little grass, *Triumfetta* and *Vigna* form sparse ground cover. Some *Convolvulus*, occasional mats of *Cassipoupa* occasional *Panicum*. Mound seedlings.

This scrub, both zones, continues for some distance but gets somewhat lower and sparser, ground cover more irregular, more bare

sheet gravel. Seedlings of any kind rather scarce. A few of *Pipturus*.

One *Terminalia* bush. Around old battered but sprouting clump of *Barringtonia* are abundant *Barringtonia* seedlings, toward seaward side. Scrub gets denser again, but ground cover less.

Then a badly scored section little ground cover, but open *Tournefortia* *Pipturus* scrub 1-3 m tall. One *Tacca* plant, one *Aplousia* ridges, on gravel sheet in openings, a few *Justicia* seedlings.

Then an open area with very sparse and irregular scrub 2 m tall, mostly *Tournefortia*, with almost complete ground cover of *Thuarea*, *Triumfetta*, some *Wedelia*. The same almost across the islet except for a strip of open small *Convolvulus* just back of seaward beach ridge.

Southward the scrub gets lower, 1-2 m tall. *Triumfetta* becomes dominant

in ground cover, with some *Fimbristylis*.

Southward scrub becomes lower and denser, in places almost closed, *Scaevola* more abundant but with *Tournefortia*. Wide gravel sheet seaward with scattered very small *Scaevola* and *Tournefortia*. Almost no ground cover except scattered thin *Triumfetta* mats.

Then sparse, taller *Tournefortia* scrub with *Thuarea* and *Lepturus* ground cover. Some bare gravel sheet. Some *Vigna* along lagoon. Some *Triumfetta* and *Wedelia* inland, also.

Then mixed ground cover of *Thuarea*, *Triumfetta*, *Euphorbia chamaecrista*.

Fimbristylis, a little *Polypodium*, *Casuarina* - where ground is sandy gravel. *Triumfetta* locally dominant.

2 Crows + 1 ternature flying around together.

On ocean side the very sparse ~~of~~ low scrub continues.

Density and height inland varies. Up to 3-4 m. locally, in places closed, others open to sparse.

On lagoon side a small depression, sand bottom, no mangrove, no veg. except a couple of small mats of *Lepturus* on sides.

Then southward an area of grass with few shrubs, back of it and southward tall scrub tangled with luxuriant *Wedelia*.

Open scrub with little ground cover seaward becoming almost closed southward, then somewhat open again, on very coarse loose boulder ~~and~~ - cobble gravel.

Thin ground cover of *Lepturus*, *Triumfetta*, a none coarse substrate barely general. Much *Lepturus* in open parts of scrub. Locally dense scrub and *Wedelia*.

Scrub thick and tall, 3 m. *Tournefortia* + *Scaevola*, with *Wedelia*, toward lagoon. Then battered mangrove swamp, with *Pemphis* around edges, some tall *Bougainvillea*, 6 m. some 1 m. *Bougainvillea*, some open depression with much debris. Southward large area of debris-filled open depression, some of it with dense stand of small *Bougainvillea* saplings. *Pemphis* irregularly peripheral.

Southward another smaller depression with *Bougainvillea* saplings, some *Pemphis*, some open ~~with~~ gravel & debris bottom.

They surrounded mostly by a very dense tall scrub of mixed composition *Pisonia*, *Allophylus*, *Pemphis*, *Scaevola*, *Wedelia* etc. Extreme end is a gravel flat partly covered by a dense & open low scrub, of

Tournefortia, *Pemphis*, *Scaevola*, tangled with *Sporobolus*-*Laba* grass *Pemphis* + *Guettarda* ~~very~~ ^{badly} beaten by typhoon but sprouting abundantly choked with *Wedelia* in point.

Seaward a dense & open scrub of *Scaevola* and *Tournefortia* 1-2 m. tall

Many large white terns flying over this area.

The scrub types on this island mostly give the impression of being even-aged stands, regardless of density, and with remarkably few seedlings in openings. Little correlation with substratum, only broad correlation with place on island. Generally sparser and lower toward seaward side.

1960 Marshall Is.

Oct. 23 - north end of
Mejatto I.on open bare gravel
sheet from 1958 typhoon4/24/14 *Fimbristylis cymosa* R. Br.
common2' { 15 *Lepturus repens* var. *subulatus* Forst.2' { 16 *Lepturus repens* var. *septentrionalis* Forst.2' { 17 *Lepturus repens* var. *septentrionalis* Forst.14-1 18 *Boerhavia tetrand* Forst.
rare

2 19 moss

on old rotting coconut trunk
in scrub back of beach.

Coconut planting on
Jaluit atoll in form of a
20' grid over most islets,
and generally regardless
of soil and terrain - swamps,
rocks, gravel, etc., and ~~is~~
regardless of existence of
standing trees. Three
foot wide and deep square
holes dug - ~~traces~~ - vegetable
trash thrown in, then hole
nearly filled with sand
or gravel. Coconuts planted
later in these holes, not
completely buried.

One rather thorough
brushing out done at times
holes are dug. Trash allowed

roots fragrant
with peppery fragrance.
tufted

(but toward
var. *subulatus*)

prostrate, flowers
bright pink.

To rot on ground, very
little evidence of burning
seen. Addition of humus
to soil is notable.

In general, exposed
gravel sheets are dark
gray or almost black
from algae.

general notes -

Lizard populations seemed large on most islets visited except Likon and Lijeron, where only a very few lizards were seen. Most seen generally were the lined skink. Black skinks were seen on Mejuinak and Jaluit, and slender skinks on ~~Jaluit~~ Kinajong and Mejatto, no each. However, no special effort was made to examine lizard populations.

Rats were seen in abundance on Enybon, all apparently *R. exulans*. Rats around the radio station on Jalut were a larger rat, but some smaller ones were there, too, judging by reports of Knudsen and from abundant droppings seen.

Cats abundant on Enybon, seen also on Jaluit. A few dogs generally, also chickens. (no dogs on Jalut, said to be several others on atoll.)

A pet Micronesian fruit dove on Enybon.

Wedelia leaves rather generally somewhat eaten by noctuid larvae (similar to army worms) but ^{plants} by no means defoliated.

Hypolimnas common.

Various ants common.

No striking abundance of insects anywhere.

Aedes occasional on all islets, but not abundant. Some Culex on Jalut, not many.

Very few insects came to lights.

Little soft wingless crickets common in Pandanus leaves on Mejatto.

House flies and a smaller similar fly occasional, annoying but not at all abundant.

Large land crab holes prominent on most islets, incl. Lijeron, where the land crabs mix sand with the humus in places much as the shearwaters do on Heron. (these holes are presumably those of land crabs, but maybe coconut crabs or even shearwaters (but a bit small).)

Land hermit crabs not specially abundant.

Data on breadfruit disease on Pundap Islet from report by Eric Knudson based on information gathered May 5, 6, 7, 1960.

	seedless	seeded
Total trees living	307	73
" dead trees	134	
" diseased trees	24	
Trees died last 3 mo.	20	4
Trees diseased last 3 mo.	10	0

Two areas of infection and a third developing - in the latter ~~the~~ tree is in last stages before leaves drop off and surrounding trees are beginning to turn yellow.

site #	dead	yellow
1	1	4
2	2	3
3	0	4

Symptoms

Tree becomes progressively chlorotic; approximately four months after symptoms are obvious lvs. fall rapidly (except for half dozen that turn green suddenly and then almost immediately yellow and fall - not personally observed).

and, as no new growth appears, roset, etc. the tree is presumed dead.

leaves: 1 top leaf - light green, yellowing starting from leaf margin, random spotting (reddish), and occasional "hairline" necrosis along the leaf margin.

2 leaf necrosis more in evidence, leaf tips slightly curled, necrotic spotting, and definite yellowing over-all.

3. increased necrosis and darker yellowing

4 similar to 3 but leaf begin to die rapidly.

Roots: A tree in the final stages before death was completely dug up and pulled down. Tertiary roots immediately underneath the trunk and those roots

5-15 mm in diameter showed the following deformities:

1 Root was twisted, wrinkled, and often constricted.

2 Galls were evident - approximately twice size of root diam.

3 These galls appear to be "stung" (a black dot from which

streaks radiated). The root xylem on either side of the gall usually had a streaked appearance, discolored black.

4 The galls themselves had signs of rot within the tissue. In the galls the xylem remained unaffected but the surrounding tissues were darkly discolored.

5 In stripping the roots black streaks were randomly but consistently found.

6 Where root was constricted the xylem remained relatively unaffected.

Surface roots within 10 ft of the trunk appeared normal. Small holes just to expose the surface roots were dug around several trees to a depth of 1-1.5 ft. There was a relationship bet. the degree of yellowing and the occurrence of bearded roots. Also a greater number of roots were found to be abnormal near the path than were found at an equal radius opposite the path.

Trunk: The tree was sawed in three places:

at base, at middle, and near top. The xylem and phloem appeared normal except for a definite black "shadow" zoning, but penetrating the xylem tissues. This discoloration had a fairly definite border in the xylem but in the phloem it slowly faded without any clear limit. This condition was found only in the tree trunk.

Branches appeared normal.

Summary

There appears to be sufficient evidence for a conclusion:

- 1 The majority of the diseased trees are situated along the path.
- 2 The disease is spreading most rapidly along the path.
- 3 The roots nearest the path show malformation before those farther away.
- 4 Once a tree yellows and dies in a recognized manner those trees nearby tend to display identical symptoms.
- 5 The time intervals between yellowing and

death, yellowing and subsequent infection of nearby trees, or death and subsequent yellowing of nearby trees, are uniform.

v. The trees show symptoms easily recognized and can be attributed to a common factor.

[this far from report]

The tree that I looked at that was affected was sprouting a bit from base, upper parts dead, but Knudsen says this is not generally the case. Surrounding smaller trees somewhat chlorotic.

When asked if some coconut trees still retaining leaves after typhoon had died later several Marshallese replied in affirmative. gave different periods of time, from 4 months to two years.

"Laure" - poor coconuts.

Neil Morris, agriculturist on Yap, studied this breadfruit disease further.

Oct. 29 - plane trip Jaluit to Majuro - ~~including Jaluit~~ ^{up and down s.e. reef of Jaluit} ~~but~~
Down Long Islet - surge channels abundant

On Jaluit - swamps only in east end.

Up Long Islet
Jaluit - photos
Pass - 2 photos

st. channel w/ bifurcations and an "island" reef in channel.

strip ~~of~~ ^{with} from Enybor - ~~remains~~ ^{remains} about recovered. Beyond this near angle in reef ~~most~~ ^{many} denuded areas scarcely recovered.

All along the lagoon reef flat bars are a very frequent feature, mostly near the ~~outer~~ lagoon edge, but also many small arcuate ones near shore. Many remnants of consolidated material, also, out on flat in various positions.

Majuro - ~~s.e. corner~~ ^{s.e. corner} lagoon reef flat ^{is wider} than seaward side. on north east and north sides

The seaward flat is much wider. Margins of seaward flat on s.e. corner more lobed. That of lagoon reef more scalloped on north side.

Almost no coral patches or pinnacles in east end of Majuro lagoon.

^{SAMOU} *Sambucus* ^{longifolia} plant, ~~in adm.~~ ^{in adm.} islet. Asked Mackenzie to send me specimen. (sent & received later)

Wedelia seems to be generally the most abundant plant. *Pluchea odorata* has become relatively scarce except along air strip, where it is still abundant.

Many ornamentals in administrative town.

1st 19th photos on Kod. roll L are aerial shots along north and west reefs of Majuro climbing from 4 to 600 ft. (last shot of coral patches in west end.)

Oct. 31 - Kwajalein Islet, Kwajalein Islet

41420 *Dentella repens* Forst.
5 common in weedy lawn near school building constructed

Oct. 29. Jabwot - flew over very low several times.

Scaevola series

Calophyllum anophyllum

Tournefortia argentea

Thunbergia innotata?

Centropus altissimus (s.l.)

Cross mucifera

Carica papaya

Quettarda & *speciosa*?

Low generally planted to coconuts, some ^{young} patches of breadfruit. (several dead and some yellow breadfruit trees some dead *Tournefortia* back of beach fringe on one side.)

Fairly broad strip of native vegetation on one side (east?), narrow on others. A large taro pit near west end, extending almost across island.

This apparently has *Cyrtosperma*, but does not look well tended. There are several houses and people.

Took about 13 Kodachromes.

Breadfruit groves mostly present a very dense and even level canopy.

prostrate, forming dense mats, flowers white.

Marshall Is. Nanna-Koraj -
Wale

Oct. 29 - Nanna atoll
flew by at a little distance
except very near south
end, only one tiny islet
on entire west reef. Many
on east reef, well distributed
except for a considerable
expansion of reef near north
end that has only 2 tiny
specks of land. The
largest islets are at
the north and south points
and on a loop to the east
in the east reef.

A little surf is breaking
along entire east reef, none
on west reef.

Oct. 29 - Kwajalein
Watched cat playing with
a mouse in front of hotel.

Mr. Boyd, U.S. Weather Bureau,
says that when he left Wale
2 years ago there were 50-100
wild cats in the woods
on Wale; a few rats at that time.

He says also that the various
weather stations, such as Kwaj, Eniwetok,
Majuro, Johnston, etc. which have
recently been taken over by the
Weather Bureau now publish
monthly and annual summaries.
Also there is a Pacific summary.

Kwajalein

Look up
Crowl, P. A. + Love, E. G.
Seizure of the Gilbert and Marshall
1-4/4, Washington, 1955.

Oct. 30 - Kwajalein - beach
and edge of reef at rather
low tide

13 Turnstones
1 whimbrel
3 plovers

Oct. 31 - trip in SAR
 albatross plane around
 Kwajalein atoll
 left field 8:40

Ebeye - most of trees
 gone from north end,
 many coconut trees
 along lagoon shore of
 settled part.

Center part of ^{Bakeri} ~~Rogee~~
 cleared, two ends well
 wooded. Construction

^{oicci}
 Bigej mostly well wooded
 but the old ramp has
 no woody vegetation,
 thoroughly green with
 Wedelia or Ipomoea.

Small islet n. of
 first pass has forest
 but no coconuts. Also nest
 on well up reef.

Reef ^{flat} finely and closely
 striated transversally, some
 channels more coarsely striated.

Next small islet also has
 natural scrub, no coconuts.
 (Also the nest (prob. P. ^{harris}?)
 also nest.)

Nest is large, some coconuts
 on south half, north half
 scrubby forest.

Nest islet scrub forest.

Nest scrub forest with
 a few coconuts.

Nest, tiny one, very few coconuts,
 scrub.

Nest one scrub forest with
 much Pandanus.

Nest scrub.

Nest coconuts.

Nest scrub.

Reef: ^{Namur} scattered
 coconuts, denser in places. transverse
 air strip usable. Much
 open ground on other old
 strips, grassy with scattered
 brush.

Nest islet west from coconuts.
 Then another with coconuts.
 Then a pass with a bare
 sand bar inside it about
~~500 m.~~ 500 m. or more.

Then a small densely
 wooded islet.

Large islet with coconuts.

Then a pass with a good
 bifurcating channel
 w/ bifurcation on it. 3 on left.

(photos but not from good
 positions - end of roll)

Small scraps of denuded reef platform west of pass.

Small islet, densely wooded.

Another, *Bisemia* forest with fringe of coconuts along inner beach.

Next, solid forest, wind-sheared on east side.

Shallow pass.

Small wooded islet.

Barf on both sides of reef, less inside.

Larger islet, with considerable coconut plantations.

Cross striations on much of the reef here.

Range islet about 20 planted to coconuts, some forest on seaward side, some of plantation extending to sea side.

Shallow wide gap.

Then reef with small bowl-shaped black gravel bar - possibly consolidated.

Small forested islet.

Islet with irregular forest and a few coconuts.

Then another with about half in coconuts.

Small bar with a dozen coconuts, no other plants.

Small islet, inner half open scrub, then belt of coconuts, then narrower outer band in forest.

Many patches reefs.

Then large reef flat area.

Islet with open scrub on east end and lagoon shore, coconuts on outer side. Then 2 small islets with much bare ground & little scrub.

Then larger islet mostly open two patches of coconuts. Small islet outside this with ^{linear} coconuts - connected by linear bar.

Small islet with open scrub. Linear bar to outside of reef.

Then large westernmost islet, irregularly planted to coconuts, a band of sand with open scrub ground south and west side. Low forest in center.

Reef much broader than islets in this western prolongation. Islets look as though a typhoon had swept them except the end one.

Marshall Is.

Back from end a number of lines, bar just before lagoon begins. One, at least, seems to be beachrock, others scarcely.

On much of northwest reef before west end there seems to be a small development of an algal ridge on the inside with some surf breaking over it.

Back along south reef several strips of reef, shallow pass with broad transverse striations.

Islet partly planted to coconuts some forest, some very yellow scrub.

Then wide gap with shallow areas and 3 deep but narrow cross channels. This reef, well submerged, is quite complex, but does not seem to have surge channels. Has sandy areas at different depths. Then a deep channel with

Kwajalein

large V sloped reef inside, each fork bifurcates. Inside of right fork a number of patches not forming a very good pattern. None of this would be awash at lowest tides, probably. Well submerged.

Then a shallower stretch with a patch of surf. Then a narrow deep channel then more well submerged reef.

Then a ~~to~~ rather long islet with loose *Pisonia* forest and a few patches of coconut trees, some open scrub on s.e.

Then a small islet with open scrub, a few coconuts, a sand bar with a few coconuts and bushes.

A wooded small islet.

A long islet with coconuts and a long spit extending lagoonward along a deep bifurcated channel.

A coconut planted islet with a long spit extending east along reef.

Another long coconut planted islet. A deep pass with

an islet inside,
very sandy, with
coconut trees.

Then a long narrow
planted islet with
a narrow brushy eastward
extension with a ~~road~~
spit in it.

Then a long islet,
irregularly planted
to coconuts, a partially
drying reef, another islet
w. half in coconuts,
east wooded and attenuating
to a spit. the brushy.

Then a very long narrow
islet with a spit westward
a deep pass with three
islets inside, it arranged
in a bifurcating
system, inner one esp. ^{sandy}
sandy. Coconut trees.

Then a long islet
the inner west part
planted to coconuts, very
elongate eastern part
in sand, one or two
patches of coconuts, then
almost a gap, then more
spit and an enlargement
with coconuts. This
irregular interrupted
sand ^{partly}
in sand. Then a small

islet, wooded with
some coconut.

A shallow pass,
sand bar on reef
no vegetation.

Then an islet planted
to coconuts.

a deep pass with
a reef with sand bar
inside.

Then an islet with
coconuts, narrowing
eastward, the part
brushy.

Scattered bar sand
or gravel bar on reef.

A patch of worn down
consolidated platform
on bar with 2 coconuts
inner bar.

Some large patch reefs
inside, but no passes
into the very shallow
reef. Elongate narrow
bars and much
dark rubble or consolidated
platforms.

Then a wooded islet
with a hook on inside
w. part end, then a
wide pass.
Large islet, densely
planted to coconuts
except for a couple of spots

inside with large trees
and opening
Another islet with
hook on e. end, and
a deep pass

An islet planted to
coconuts, narrowed
eastward to point.

Pass on reef, one with
a coconut tree.

Small irregular islet
a stabilized bar with
Scaevola. Complicated
system of cross
bars, then an islet
with coconut plantation
then a deep channel,
with a wide gap,
shallower eastward.

An islet mostly in
coconuts, some *Pisonia*
fruits some scrub, mostly
scrub on east half and
a patch of *Templum* on
east end, with depression
inside vegetated high
gravel or sand ridge
at pass. Wide deep
pass. Then a couple
of reefs with small
bars.

An islet with ^{tall}
Pisonia, some coconuts
connected by a ^{small} *brushy*

bar with another with
a hook on east end.

A unadorned bar
laggedward of these two
islets, almost the length of them, ^{wide,}
^{separated from}

Then a wide ^{deep}
pass and gap with
submerged reefs.

Then an unvegetated
partly blackened bar.
A couple of large
patches inside with
bars on them, no submerged.
A tiny islet with coconuts
a deep narrow pass.
then an elongate islet,
irregular width, planted
to coconuts on wider
parts.

Then a shallow pass.
then a narrow elongate
islet or bar with
some coconuts, a little scrub.
A small bare bar. Then
a shallow pass.
A reef with an elongate
narrow bar, widening
eastward into an islet
with a well pointing hooked
bar on west end in lagoon,
narrows to bar on east.
Shallow pass.

Elongate islet, west part
partly bare, partly ponds.

partly in *Pemplicis*
 eastern part wooded.
 Deep pass, bifurcated.
 Islet with *Pemplicis*
 surrounded pond,
 some coconuts.

Then an elongate islet
 partly in coconuts,
 somewhat beaten up,
 eastern end wooded.
 Then a rather deep gap.
 Then a wide reef
 with ^{very} narrow elongate
 very narrow ^{irregular}
 bars, interrupted
 anywhere.

Then a large islet
 open, scattered Pandanus
 & other trees mostly
 a blanket of *Wedelia*
 scattered coconut trees.

Then shallow gap
 to Kwajalein.

Arnold LC 7 on reef.
 Went on north horn
 Kwaj. 2 bare islets
 or bars, two tiny islets
 with scrub.

Everywhere many coconuts
 in south part of village
 and around C.G. station
 none on north half.

S. & N. Loi open, beaten
 up. Some coconuts some

brushy low forest on
 North Loi, elongate and
 brushy ~~and~~ northward.
 Small brushy islet
 stepped beach rock
 along reef, then very
 long narrow brushy
 islet, attenuate
 westward.

Then Baker (?) with
 considerable construction
 in middle bar section.
 Then a large islet,
 coconuts in east 1/3,
 mostly in low dense
 forest.

Then a moderately
 deep pass. Then Bigaj
 Ruined tanks still there
 with vegetation around
 them. Coconuts on w. side.
 Long reef. Projections
 of living coral and algae
 extending into lagoon.

Next islet in coconuts.
 Islet in pass Kwadath,
 mostly forested, a few
 coconuts.

Cinivatah still dense picnic
 forest with lot of bird life
 but e. end toward pass
 has a grove of coconut.
 Some rock on this point
 awash.

Northward on reef are some curious circular reefs extending from main reef into lagoon, active growth around periphery of these, sand in center.

Flew south to Lib J.

Reef is widest & sweet. Pond in n.e. corner. N. w. corner and n. side has considerable forest, *Panicum*: with only a few coconuts.

Forest around pond not *P. amplius*?

Patches of breadfruit in

Tournefortia *Calophyllum*
Scaevola *Pandanus*
Antiaris *Bruguiera*
Hibiscus etc.

Wind grooves on n. side diagonal to beach.

Surge channels on n. & s. sides, scarcely on west.

The forest around the pond is puzzling - does not look too much like *Panicum*, nor *Bruguiera*, one patch may be *Bruguiera*, brighter green than rest. Pull green, rather small

righter

leaved, somewhat tessellate canopy.

The pond is brown, rusty looking.

This island should repay study.

Back to Kwajalein,

North along north reef ^{at very top of} point of point Baber Bigej etc.

Islet bet Baber & Bigej has very considerable *Panicum* ^{complex}

Extensive boulder tract on reef north of Bigej. Boulders tend to be aligned transverse to reef, accounting for some of striate appearance.

Most of dark patches on this reef are these boulder or rubble tracts.

North of Kwadab - a huge loosened boulder - a part of a butters between surge channels, lying ready to be lifted on reef by a storm.

Coarsely striate appearance apparently due to submerged surge channels.

Rubble all along shows a remarkable tendency to transverse alignment. But fine striation on paler parts of reef flat is apparently not due to this, perhaps to shallow grooving such as seen at Kwaj by terminal.

Most of forest seen on these islets earlier this morning is *Pisonia*.

The herbaceous vegetation on the old ramp at Bigaj seems to be *Wedelia* on about the peripheral half or $\frac{2}{3}$ and *Sporoclea* in center, though hard to be sure.

^{1.46} A very few scattered patches reefs in lagoon. Lagoon is quite rough today, quite a few white-caps.

Down south reef at low altitude

Little or no alignment of rubble on these reefs, bars very numerous, though annular bars on lagoon reef flats opposite islets are occasional.

Pamphus abundant occasionally showing wind striation.

Oct. 16 - Honolulu

41421

Cinnamomum bakeri Engl.
cultivated in garden

Oct. 25 - Honolulu

41422

Cinnamomum asiaticum L.
cultivated
(brought in by C. M. Oliveira)Nov. 4 - Trip by Boeing jet
Honolulu to San Francisco

10:30 a.m. - 6 p.m. 34,000'

overall that
 much of the
 was observed
 from a
 slight
 upward

At about 1:30 - 2 p.m. Honolulu
 time the sea, as such, was
 practically unseeable - a
 scattered layer of small
 clouds ~~was~~ lay far below,
 spotting what appeared
 to be a deep blue sky, slight
 duller and richer than the
 "iodochrome blue" seen in
 high mountains, but
 neither surface of sea nor
 horizon visible. The clouds
 did not differ in appearance
 from what they would have
 looked like in the sky above,
 except that one looked down
 on them. A slight haze
 obscured where the horizon
 would have been, fading

from seeds brought from
 Utaite in 1956 (Fosberg 36713)
 Flowers picked by C. M. Oliveira, ^{earlier}
 from same infl. as collected
 dry now.

out both upward and
 downward. Above the
 sky was almost a
 blackish blue - not a
 night sky, no stars, but
 simply a sky blue
 diluted with black,
 a rather opaque blue,
 not transparent, lightening
 to a normal sky blue
 toward where the horizon
 should be. Being on the north
 side of the plane the place of
 the sun in this color system
 was not evident. From
 time to time an invisible
 turbulence shook the plane,
 giving a rather eerie effect.
 (As near as I can see from
 the other side, an indirect view
 through the glass diagonally, the
 sun is just a intense white light
 scarcely influencing more than the adjacent blue.

Nov. 5 - Miles Canyon

still in fairly undisturbed condition except for railroads and highway in bottom.

South wall mostly densely wooded with a low forest of live oak, *Aesculus*, *Arbutus*, etc. Other wall open, grassy with scattered live oak. Stream in bottom with abundant water plants.

Road to Sunol, more open, but scattered live oaks and *Quercus lobata* grass heavily grazed.

Nov. 7 - Sacramento Valley,

from air. - The area west of the lower Sacramento River is almost entirely under intensive rice cultivation. The land is in large blocks and the entire area is cut by a system of large sloughs lined by very narrow gallery woods, usually not much more than a single row of trees, or even only a strip of low scrub.

Between these sloughs are a network of ditches. Along the river, itself, there is a narrow to broad

belt of orchard (peaches?), mostly on the east side, still green but showing some change of color toward red. At a junction of several sloughs is a patch of forest or swamp. Some small wooded islets in larger sloughs.

The soil, where the fields are ploughed, is almost black.

Nov. 22 - Chantilly Airport
Large open field, thickly
grown with weeds.

Different weed communities
apparently representing
different times of withdrawal
from cultivation - boundaries
between them abrupt and
straight. One community
dominated by *Bidens* sp.

A few scattered *Rubus*
and one *Rhus* in this.

Separated from a
grass community
by a line of *Juniperus*
1-2 m. tall, the *Bidens*
spread 1-2 m. beyond
this into grass. *Bidens*
apparently annual.

In smaller numbers
on into the grass.

In the *Bidens* abundant
small burrows into
ground - as many as 7 in
one sq. m. seen.

In grass a maze of
intersecting runways,
some burrows. Also
some larger burrows
with fresh dirt
around mouths.

2 or 3 *Juniperus* seedling
up to 6 m. tall in grass.
A few scattered small

clumps of *Andropogon*.

Several rows of large
Juniperus.

At least 6 or 8 ^{more} hawks
of at least 2 species flying
mostly just over the tops
of the weeds and bushes.
Several meadow larks.
2 shrikes.

Remains of mice, etc.
occasional in grass.

Raccoon tracks in mud
near large puddle.

One large owl just at sunset.
also flying low over grass.

One row of oaks with
a small patch of saplings
nearby.

Cytisus
types

Tripp Run arm of
Lake Barcroft
Jan. 22, 1.5 mi. s. of Falls Church

edge of deciduous woods,
shore of lake.

- 41423 *Betula nigra*
occasional
- 2 24 *Ligustrum ligustrinum*
occasional
- 2 25 *Vaccinium*
occasional
- 5 26 *Alnus serrulata* Willd.
common
- 5 27 *Corylus*
occasional
- 2 28 *Famnamelis virginiana*
occasional

After several days of weather ranging between 25° and 0° F the lake is frozen over with a firm sheet of ice, that bears a grown person's weight very well. There are several long irregular cracks that have frozen together again, after opening. On the surface are patches of ripple marks, also curious series of concentric irregular ring-like ridges, also patches of semi-consolidated snow.

The upper part of this arm of the lake, the down

small tree ^{ca. 12 m. tall,} with rough bark; winter condition.
shrub 1 m. tall; winter condition.

shrub 1.5 m. tall; winter condition.

many-stemmed shrub 2.5 m. tall, catkins pendent, not yet at anthesis.

shrub 1.5 m. tall, catkins pendent, not yet at anthesis.
shrub 1.5 m. tall; winter condition.

mouth of Tripps run, is largely filled with silt and covered by marsh. The lower parts of this are dominated by tussocks of *Juncus effusus*, the upper and greater parts ~~with~~ by *Typha*, with clumps of *Sagittaria arifolia* (red aff.).

These marshes do not seem nearly as firmly frozen as the open water. The ice here, as well as in muddy places around the edges, seems mushy. Possibly there is a slight elevation in temperature from heat from decomposition of organic matter in the mud.

Feb. 22. A heavily overcast day, damp and locally very foggy.

In Washington, in the afternoon, areas where snow was still lying were observed to be apparently smoking - white vapor was drifting up from them in great quantities.

In the evening, during considerable driving around the area south of Falls Church and around and southeast of Seven Corners, fog was locally so thick as to make driving almost impossible.

The distribution of the fog was very irregular. It was observed to be much heavier in areas where there was still much snow on the ground. The snow was "giving off" clouds of vapor, and where it was piled up the snow piles looked like smoking volcanoes. The temperature was generally warm,

probably at least in the 40's. Obviously the fog resulted from warm saturated air drifting over the snow and being cooled off, forming clouds, and the irregularity of its distribution resulted from the irregularity of the persisting snow from storms a week or more earlier.

Feb. 26. Great Falls -

The water is so high that only one small patch of rocks is showing. The water is a cafe au lait color and smells quite filthy.

Mar. 3 - Route from U of H to
Wilson Tunnel to Kaneohe &
Lanikai.

Delonix regia along Wilder Ave.
Foster Garden. *Spathodea*
on corner.

Large *Samanea* in Palama
Mango trees. *Cycas* at
Kupahala St.

Kam School on mountain
to right. *Casuarina* forest
above it and on ridges to
it. • left of Kalihiki St. on
way to Tunnel.

Center row of *Jasminum*
multiflorum between highway
Below *Casuarina* on slope
scrub largely of *Schinus*,
more *Psidium* and *Eugenia*
cuvini up the valley.

Scattered *Acacia* *koa*, abundant
more abundant up the valley.
Patches of planted *Eucalyptus*
Eugenia jambos, then
forest of *Melaleuca*,
Casuarina, etc. just
below tunnel.

View of Kaneohe Bay.
Plated cliffs with
Pandanus forest at base
at left. *Gleichenia* patches
Banana plantations. Large
mango trees. • Road
lined with banana

plantations down to
Kaneohe.

Large mango trees and
Spondias? on rt. before
main intersection.

(Kohohahi - Kailua Rd.)

Hills on right with
Lantana, *Leucaena*, *quava*.

lowlands on right mixed
low forest of *Hibiscus tiliaceus*,
Eugenia cuvini, *Samanea*
quava, *Leucaena*

Prosopis forest at base,
low, flat-topped, apparently
wind-sheared, undergrowth
of *Schinus*, etc.

Mar. 4 - University to Moku-
leia.

Wilder Ave. -

Punahou - *Hylocereus* hedge
Row of *Rousettea*

Kunahilo St. bet Ward

& Victoria - fine *Ficus*
microphylla (a *benjamina*)
on rt. in yard.

Vineyard at Queen Emma

fine groups of palms.

Vineyard & ~~Houghtaling~~ Palama
on left a *Phoenix*, prob.

hybrid of *P. canariensis* & *dactyl.*
2 large *Samanea* at Houghtaling.
to rt. on King, on left several
more *Phoenix*.

Large Ficus bengalensis
on King beyond Kalibi.

Moanahanu Gardens.

Salt Lake Hill - Kiawe
forest, Leucaena under layer
some Opuntia on left.

Red Hill - mostly Leucaena
scrub, scattered low Prosopis.

Area - Cassia fistula
and Acalypha hedge on it.
as road enters town

Several Cassia fistula in town
sugar cane fields.

Leucaena with some Prosopis
in waste places around
P.H., but mostly cane.

Waipahu

10597 Kohofield Rd. Jct.

Sugar cane.

Waianae Mts. ahead to it.

Lower slopes grassy and
eroded in places, mostly
thin kiawe forest, with
cactus + shrubs + grass.

10598 Barber Pt. Jct.

Sugar.

Very good Prosopis forest, about 5m tall
with Acacia farnesiana
interesting dry land plants

Sida fallax

Ebrtilon incanum

Chloris inflata very abundant

on very rocky ground.

Gossypium tomentosum toward
Nanakuli

3 bluffs on right with
thin Prosopis much cactus.

In Nanakuli large
Calotropis gigantea (white
blossom)

Bet. Nanakuli + Waianae
steep bluffs showing
lamination of lava.

ledges with grasses, etc.
Waianae - tidal flats
with Batis.

Much denser Prosopis forest.
Cliffs with cactus.

Makaha - native forest
and Moku visible at
head of valley.

Kiawe forest on debris & talus
slopes

cactus, pili grass on ledges.

Flats beyond - Heave
forest, fractured.

herbally - quartz rock slope.

Vitex on sand at top
of beach.

Cane.

Kiawe forest

Makaha V.

~~Kiawe~~ Kiawe forest.

Kaena Pt. road. at base
of cliff. Kiawe scrub
+ pili

Strand flat, with
Sporobolus.

Lower tuff slopes
with *Atriplex semibaccata*
Sida, *Trinachne*
Jacquemontia, *Chloris inflata*
Heteropogon (abundant)

~~Small bushes with~~
Euphorbia.

Santalum *Waltheria*

Chenopodium

Boerhavia

Low *Leucaena* shrubs

also *Acacia farnesiana*

On sand dunes

Sesuvium

Sesbania

Heliotropium

Euphorbia degeneri

Boerhavia diffusa

Atriplex semibaccata

Dactyloctenium

Sporobolus virginicus

Vitex simplicifolia

Jacquemontia

Sida fallax

~ *Pisonia per-caprae*

~ *Leucaena*

sea stand, probably eolianite
or consolidated dune. These
remnants occur at various
elevations, up to base of present dune
(samples)

Toward Mokuiaia

Leucaena scrub

Gane fields and
gasuarina wind breaks
Prosopis - open stands
with *Leucaena* scrubs.

The slopes around
Kaena Point are mostly
covered by a *Leucaena*
scrub that is dead in
its upper parts, possibly
killed by salt spray,
but with prostrate
branches at the bases
of the dead bushes
that are living and
green. The whole vegetation
seems to be in poor
condition (because of
dryness or salt?).

On the windward side of
the point is a great
mass of lava boulders
piled out over a rough
terrace just above high
tide. On this are remnant
patches of what appears
to be beach-rock - bedded
lime sand and gravel, quite
case hardened. If beach-rock
it is from a previous high

Mar. 6 - Univ. To Koko Head.
 Leucaena scrub on hills
 to left of road below Univ. far
 beyond Kaimuki - Kapakahi
 Valley - lower part the
 valley is filled with
 coral plain. Lower
 slopes covered by scrubby
 open Prosopis forest. Coral
 plain built up, golf course
 + residence area. Formerly
 cane field. Double highway
 with Jasminum multiflorum
 and Lantana sellowiana
 in middle. Hibiscus hedges,
 also Nerium hedges.

Wailupe Valley - some
 native forest and Kuhlmannia at
 head, slopes Prosopis
 with considerable pili
 grass. Some Opuntia.

Beyond - a row of Delonix
 on left. Bougainvillea
 on bluff at right.

Nin Valley - Kuhlmannia very
 noticeable at head, slopes
 on side of scrub (Leucaena?)
 with scattered Prosopis.
 Lower slopes Prosopis. Some grass
 on higher slopes.

Hibiscus tiliaceus on right.
 Kuhlmannia V. same as
 Wailupe; Nin. Ridges badly
 eroded. Goats?

Kunapo Pond - extensive
 margin of Batis maritima,
 but being destroyed.

Maunaloa Bay shallow
 reef.

Koko Head - badly eroded
 Prosopis and Leucaena.

Koko Crater - Prosopis
 on base, grass, some
 spots of Leucaena near
 top.

Hanauma - Prosopis
 on slopes with loose
 understorey of Leucaena.

Prosopis on flat rather
 dense, no undergrowth -
 cleared out for picnic ground.
 Steep slopes around
 bay with patches of
 wind-sheared Prosopis,
 interspersed with patches
 of grass and erosion scars.
 Kaula

Walk down path from parking
 lot.

Weedy veg. of Chelidonium
 Alysicarpus, Passiflora
 foetida, Waltheria Leucaena,
 Atriplex, Tricholoma
 Bluffs of bedded tuff.
 Boerhavia repens
 Aida
 at base Leucaena thicket.

planted Casuarina Cocor,
 Ficus microphylla, Malvaceae
 Polyscias quil. + Plumeria
 Canna, annual pavillia.
 Cyrtosperma
 (Weeds - Chenopodium murale,
 Cirsium, Eleusine, Chloris
 inflata.

Sporobolus indicus

Zizania glauca

Reichardia

Sida sp. *mentha* complex

Lonicera olivacea

Pennisetum verticillata

Ocimum

Dactyloctenium

Malva parviflora (on sand)
 upright, lvs. pub. above.

comp. tuft beds on it.

Merremia aegyptia

Verbena (on sand)

Tribulus (on sand)

Portulaca maritima (top of beach)

mangrove fruit on beach

Patch of *Thespesia* at
 base of slopes at end of
 beach to right.

Rolls up through it.

Lipochaeta integrifolia

Sida - (different)

Hibiscus tiliaceus (green beneath)

above -

Sporobolus cainii

Portulaca oleracea

Stachys arvensis

Ocimum (anast. l.)

Jacquemontia

Rantana

Eragrostis variabilis

Heteropogon contortus

Low scrub of *Prostrep. acacia*,
Pantana, *Tricholoma*, ^{*leucocoma*}

Passiflora, *Reichardia*

Waltheria

good patch of *Eragrostis*
 on eroded place.

Gonitis javanica (dwarfed)

serious erosion on left
 side with *Eragrostis*

Emula lonchus, *Sida*

Chrysopogon, *Portulaca* in corner
 of tuft.

Eroded trail up through
 scrub of *Acacia farnesiana*

Cassia leschenaultiana

Ageratum conyzoides (dwarfed)

Top of ridge - bare, eroded
 down to tuft layer surface -
 (dead)

Panicum faurei

Chloris, *Dactyloctenium*

Waltheria, *Heteropogon*, *Lonchus*

Heteropogon, *Atriplex*,

Reichardia, *Lipochaeta*

Passiflora, *Digitaria*,

Fimbristylis cymosa,

1961

~~Common~~ Hawaiian Is.

wind-sheared *Prosopis*
on right

Walking along Hanauana
slope toward sea end -

First a very smooth
eroded tuff, then a rubblely
tract, then a grooved or
gullied slope.

Heliotropium curassavicum

Fimbristylis, *Panicum*, *Richardsonia*
Dactyloctenium, *Lipochaete*
all in pits.

First small crater
Hanauana slope dense
grass - *Chloris*, some
Atriplex, *Cynodon*.

with ~~#~~ *Pilea*, *Richardsonia*
Emilia - *Pilea* is prostrate

Crater floor surrounded
by low wind-sheared scrub
of *Prosopis*.

Inside this more grass
with *Commelina bengalensis*

In lowest part of crater
solid sod of *Marsilea*
but with considerable
invasion of *Chloris*, *Seteria*
& *Xanthium* a little

Richardsonia, *Mammisia*
Malva, *Perinochloa*
Colocasia, *Polygonum*

Phaseolus lathyroides

Emilia javanica

Jacquemontia (silvestris?)

Waltheria, *Sonchus*

Tricadum, *Cynodon*

Paspalum foetidum

Ageratum, *Solanum nigrum*

Malvastrum

From crater, west - striking
silhouettes of wind-sheared
Prosopis on horizon.

4-5 ft of water, up to edge
of *Pilea*.

On out toward end of
Koko Head, more bare &
eroded tuff.

Around edges of crater
some remnants of ~~the~~
silty soil held by
grass, and tangled roots
of *Maave* - belt about 2 m.
from *Maave* intact, then
2 m. more eroding.

Scheuchzeria palustris scattered
on eroded slopes (typical var.)
locally common - seedlings

Seaward slope also eroded. Patches of *Prosopis* scrub *Heteropogon*
 ✓ *Chloris* in small ravines
Schinus molle common.
Waltheria, *Pennisetum*,
Aida, *Jacquemontia*, *Atriplex*,
Emilia, *Portulaca oleracea*,
Atriplex

Down slope vegetation
 disappears. Last straggling
 plants are *Chloris*, *Dactyloctenium*,
 & *Portulaca*. *Schinus*.

Panicum or small trees
 Wet with spray on
 breeze day up at least
 29-31 m.

Spectacular wind erosion
Nama sandwicensis
 rare in crevices.

Euph. hirta, *Lipochlaena*

Hanauma slope

Down slope, crumbly,
Lipochlaena, *Schinus*
Aida, *Desmodium biflorum*
Panicum Port. *oleracea*
Pennisetum *Waltheria*
Dactyloctenium
Euph. hirta, *Stachytarpheta*
samoensis, *Jacquemontia*
Panicum trivium

On sheltered aspect
Vigna marina, *Emilia*
 and much more of everything
 else. (*Heteropogon* abundant.
Borhavia rare (5 years ago
 abundant and dense)
 Spectacular wind erosion
 in depression running
 down opposite crater.

Route back

Crater on left filled
 with *Prosopis* forest.

Along ridge
Asystasia gangetica
Schinus terebinthifolius

slope -

Heteropogon (patchy)
 with wings sheared
 clumps of *Prosopis*.

From top - beautiful
 view of coral reefs in
 Hanauma Bay
 of Koko Crater,
 of serrate crest of
 dry end of Koolau
 Maunaloa Bay, fringing reef
 along road
Desmodium virgatum
Euphorbia prostrata
Crotalaria mucronata

Road from Hanalei
cut through bedded tuff
Prosopis forest. Scaevola.
hard flats with
Yucca per-capras
Heliotropium anomalum

Where road turns in, one
rob. Euphorbia left on
it.

Gossypium along fence
on right. Weedy flats
Prosopis forest on slopes
to left. Cliffs above the
have Heteropogon.

Dryopteris decora on
high ledges.
Abutilon incanum
on left just before
Makeapepe pass.

Mixed pile grass
and briars scrub.
Lipochaeta - Scaevola

Stop at base of cliffs
Cliffs with Scaevola,
Schizanthus, varying from wide
Sida, Lipochaeta
Panicum toridum linear
thin, dull
cartilag. linear
Dryopteris decora
Schizanthus seedling
abundant on ledges and
valleys.

Proerhavia repens
Panicum (lawei)
Ripochaeta integrifolia
Anemone glauca (var. lawei)
Many weeds.
incl.
Cucumis dipsacens
Solanum nigrum

Canavalia galeata var.
with pubescent lvs.
acute small sepals, no
stipels maroon fls. with
green patch and white
radiating lines from base
of standard.

On down at valley house

Low scrub of Scaevola,
Pantalan ellipticum,
Prosopis (very depressed)
Yucca indica
Y. carinica
Sida
Jacquemontia
Raietrandia
Panicum toridum
Dactyloctenium
Lipochaeta integrifolia
Fimbristylis cymosa
Heliotropium anomalum
Euphorbia degeneri
Lycium carolinianum
Curcuma sandwicensis (var. lawei)

- Atriplex
 - Digitaria hawaiiensis
 - Panicum frax
 - Pectyloctenium
 - Sporobolus virginicus
 - Tribulus cistoides
 - Boerhaavia repens + alboga
 - Sponsoa per-capitata var. trana
 - Cronopus didymus
 - Ruppia prostrata
 - Arrogallus arvensis
 - Erigeron javanicus
 - Stachys arvensis
 - Tricostema
 - Arabis
 - Menyanthes aegyptia
 - Asptaria
- on Maunaloa side low scrub
of low scrub. some hantalan
- Coccoloba
 - Adiantum
 - Senecio
 - Euphorbia

Mar. 4, 1961 - Lanikai Point,
Kailua

- 41429 Stachys arvensis
1 low cliff just above beach

Mar. 6, - Koko Head
on bare eroded tuff

- 1 30 Panicum
common locally
- 1 31 Nama sandwicensis
very rare in crevices

2 70

weak herb, colorless.

60 m

prostrate

70 m

flowers purple

170

1961 Hawaiian Is.

- 41432 *Portulaca oleracea* L.
occasional
- 1 33 *Digitaria*
rare
- 1 34 *Emilia javanica*
common
- (? 35)
in ^{the} silty [&] bottom of crater
- 1 36 *Marsilea villosa*
abundant
- 1 37 *Euphorbia*
common on ^{road} roadside

March 5 - flats northwest
of Makapuu Head, on
windward side

- 1 38 *Coccoloba*
common among lava rocks

March 7 - Manoa Cliff Trail

- 1 39 (bamboo)
forming a dense brake

March 8 - Ridge north of
Opaecula Gulch, Koolau Mts.
open wet savanna-like

- 1 40 *Rhynchospora*
occasional
- 1 41 *Arthrosteremma latifolium* A. N. S.
very common
- 1 42 *Sacciolepis contracta*
common on bare cut banks
above trail.

Oahu

171

- 40 bronze, stems and leaves
fleshy.
- 60
- 60 dwarfed by unfavorable
habitat; flowers reddish.
- 100m prostrate, purple.
- 6m.
- Twining, flowers
greenish, white.
- 420m. ~~shaggy~~ culms widely
spaced, erect, 5-7cm thick
dark green, internodes
rather long.
- 425m.
- flat.
- erect, petals 4, pink,
caducous, about 1cm. across.

Mar. 7 - Mauna Cliff Trail
 entrance to trail
 row of *Cordyline terminalis*
 on left. *Psidium cattleianum*
 and *Eugenia jambos* at
Senecio sp. along trail.
Alcornoque and *Acacia* tree
 overhead.

Thicket of bamboo
 Forest of *Psidium cattleianum*
 Occasional large tree

Large patch of *Cordyline*
Psidium guajava, *Alcornoque*
Lantana along trail.
Senecio gaudichaudiana
 Trail lined & choked
 with *Setaria* and masses
 of *Eupatorium riparium*
 on banks.

Seems to be about
 on border bet. low and
 lehua zones. *Bulbui*
 patches frequent.
 Much guava forest.
 Some *Gleichenia*.

Pauoa Flats trail from
 fork -
 down hill past several
Pisonia trees, onto flat
 covered with planted
 forest of several species
 of *Ficus*, several *Eucalyptus*
Melaleuca, and *Lagerstroemia*
speciosa etc.

Large patch of bamboo.
 Then guava forest to
 pass overlooking
 Nuuanu Valley. Up
 ridge to left
Gouardia terminalis, *Senecio*
eriantha, *Pelea dusicefolia*
 (tree), several forms of
Metrosideros.

Down the draw toward
 Pauoa muddy grassland
 with *Jussiaea*, *Cramelium*
Setaria, etc. *Panicum*
purpurascens.

On slope opposite
 a very heterogeneous
 population of *Metrosideros*
 with *Wikstroemia*
sabauensis, *Senecio*
gaudichaudiana, above
 it some *S. molle*, and
 formerly, at least
 hybrids between them.

Mar. 3 - Honolulu to Pramoho
Cane fields, some Eucalyptus
plantings and pineapple fields
Kipapa Gulch.
Pineapple fields + Eucalyptus
Wahiawa. Arancaria

Gulch beyond pineapple
stand - steep sides have
some native plants -
Osterocleus, Dodonaea, leaves
~~of~~ *seicea*! Metrosideros
Mainly guava and *Leucaena*.
Panicum perpurascens.
Aleurites.

Opaerula Gulch
lower part of Metrosideros
Osterocleus, etc.

1100'

Hecht place is in an old
planting of Eucalyptus and
Cupressus macrocarpa.

Road above this through
fields with scattered lava
to sharp turn in rd. Then
small road up ridge of
end. Trail from here on
ridge.

Eucalyptus on ridge,
good lava forest on sides
of gulch, kukui in
bottom. some small lehua
at starting point in trail.
Psidium cattleianum

Osmanthus

Along trail -

some Eucalyptus planted

a *Acacia* *koa*

o *Psidium* *agrippinum*

c *Psidium* *cattleianum*

c *Stachytarpheta* *dichotoma*?

pubescent, slender spks,

obtus. acute lvs, not

sharply serrate.

la. *Melinis* *minutiflora*

o *Styphelia* *tameramea*

c *Santana* *canaria*

c-a *Santana* *geniculata*

la. *Aleurites* *linearis* (Hatch.)

o *Cassia* *leschenaultiana*

c *Metrosideros* *collina*

c *Sphenomeris* *chryseus*

c *Andropogon* *virginicus*

c *Smilax* *javana*

o *Spathoglottis* *plicata*

v *Bidens* *pilosa*

o *Psidium* *guajava*

lc *Chrysopsis* *aciculatus*

lc *Axonopus* *compressus*

lc *Saccolipis* *indica* *contracta*

o *Cuphea* *carthagenensis*

c *Santalum* *freycinetianum*

o *Osmanthus* *sandwicensis*

o *Passiflora* *edulis*

o *Aleurites* *moleucana*

o *Scaevola* *gaudichaudiana*

o (e) *Nephrolepis* *exaltata*

o *Paspalum* *orbiculare*

o lc *Diabella* *sandwicensis*

- c. *Freyrinetia arborescens*
Gibbium
Paspalum conjugatum
 2. *Euphorbia multiflora*
Erechtites hieracifolia (purple)
Cordyline terminalis
Rubus rosaeifolius
Wikstroemia sp.

Camp

flat ridge top with
Glechoma, *Lycopodium*
Cerium, scattered
Scaevola, *Metrosideros*, etc.
 some grassy spots
 with *Melastom*, *Andropogon*
Phytolobos, *Paspalum* sp.
Wikstroemia (rare)
Grevillea robusta

return

Maccharina weyereri
Psychotria mariniana
Chromolaena *Kakiana*
 side trail
Eupatorium riparium
Ageratum conyzoides
Corypha *bonariensis*

flat
 erosion
 remnant

rotte deser —

1200

park car at end of
 road on small bench
 covered by *Panicum purpur-*
ascensum surrounded
 by large acacia tree
 some small *Metrosideros*
 + *Aleurites* (tree with
 pale leaves opposite car).
 Walk along ridge
 through open acacia
 tree woodland - some
 planted *Eucalyptus* at
 first. Many weeds
 along trail. *Pteridium*
Sphaerocarpos and *Gleichenia*
 On steep slopes acacia
 tree is generally dominant
 but with scattered
Asplenium, *Psidium*
Cathartium, *Metrosideros*
 and on right small
 clumps of *Santalum*.
 A bit up the trail
Freyrinetia appears,
 growing in trees. Some
 epiphytic *Nephrolepis*,
 but very few epiphytes
 here. *Scaevola gaudich-*
audiana, *Dianella*
 appear. Local patches
 of *Cordyline terminalis*,
 and *Euphorbia* in the
 gulch bottom and

in draws on steep sides patches of Aleurites forest show up pale against the general dull green of the hooa and lehua. Bright green splashes are patches of Gleichenia locally small tough shrubs of Wikstroemia. Climb up sharp ridge to small camp.

1400-1500' Above this is an open savanna of Gleichenia and grasses, with abundant scattered Scaevola, some Metrosideros, hooa, etc.

Upper small road is through open woodland of hooa, some lehua and Scaevola - much Panicum, andropogon.

Before hairpin turn - cut with weathered yellow basalt.

View of Apanala Gulch with small Metrosideros wood on near slopes.

Quava forest and mango trees, etc in bottom, hooa on slopes, lehua in draws opposite with

hooa on ridges between some grassy slopes. ~~and~~ patches of Gleichenia opposite up stream.

Hecht play - part bet road and reservoir - flat with planted Grevillea, Cupressus, Acacia dealbata, etc.

900-1100'

Slope above reservoir has some grass, many weed. but also Coccoloba, Wikstroemia, Cordyline, Scaevola gaud. Pteridium, Styphelia, considerable Metrosideros scattered Cupressus, Grevillea, Psidium cattleianum. Some eroded slopes. Osmanthus, Osteriales

Turn ^{at} ~~at~~ ^{at} Halepa market & small church. or bet. bridges of Halepa Theater

In Area - taro patches on side of old road.

March 9 - Poamoho Trail.

Pineapple fields - then fields with *Panicum* and scattered acacia trees, many planted. *Albizia* sp., etc.

Then gulch with some native trees.

A dense stand of planted *Melaleuca*.

Then lehua forest with considerable mixture of planted *Eucalyptus*, *Albizia*, *Melaleuca*, etc. Scattered *koa* locally. *Vanilla*

stagnant. *Clidemia hirta* common along road, also

Pluchea odorata, *Metrosideros* displays usual variation.

Melaleuca common along road on ridges. *Spathoglottis* along roadside, with

Commelina diffusa, patches of *Panicum purpurascens*.

On steep bank *Gleichenia* *lyc.* common, *Cladonia*, *Andropogon*, *Sphenocarpus*, *Lehua* seedling.

Good example of this at turning place at end of rd. Trail starts through a grove of acacia trees.

Gleichenia linearis on steep bank on left.

Rubus rosaeifolius at base of bank, also *Athrostemma* *Jussiaea* (pink fls)

Small-landed *Sida*, *Gleichenia*, *Sphenocarpus*, *Nephrolepis exaltata*, *Macharina angustifolia* and *M. meyerii*. *Heavola gaudichaudiana*.

At the bottom of a small ravine a clump of bananas probably persisting from old Hawaiian times. At upper edge of this a large

Cyrtandra cf. *crispifolia*?

and *Cibotium menziesii*

Chamaecrista, *Jussiaea* along trail. Immediately

on right *Freyrieta*, *Antidesma* and *Perettia*

Then a large tree of *Bacca elatior* with

epiphytic *Claphoglossum reticulatum* and a somewhat smaller *Antidesma*

Nice Cibotium below. all within 3 m. of bananas.

Then several saplings of *Leuca* sp. and a number

of large *Metrosideros* with epiphytic *Claph. ret.* and *Alysicarpus*.

Acacia koa on slope immediately above, and

Clas carpus bifidus about 100 m. *Eugenia sandwicensis* on right.

Clidemia common along trail

- then a small fresh landslide near. Excellent examples of variation in lehua. Crossed opposite side of ridge. at crossing a clump of *Gabunia beckeri* several small Pelea, then a *Magnolia*, to left among small *Metrosideros* a few m. farther, on left a small *Des. anomala* and a *Broussaisia* (large shiny serrate lvs.) *Sphenomeris* on bank at right, and *Gleichenia* ~~thymoides~~ *glauca*. Immediately on left a *Gouldia terminalis*, *app. var. crissata*, blue berries. Then excellent *Broussaisia* and *Antidesma* and on right *Forsteria* and *Gabunia*. Across to left *Pouteria* sp. large tree coming up out of gulch, st. branched at top. *Psychotria* (*Araucaria*) cf. *mauiana* (white fls.) a large lvs just beyond to left. on it, *Gardenia remyi* several m. off trail in *Gleichenia* clump. Colony of *Gouldia terminalis* var. *crissata*

when
trail
crossed
back
on it
right

100 m.

- Then a nice view of gulch on right at curve in trail. Just about on left a small shrub of *Pilea chinensis*, and just beyond on it *Pilea* cf. *parviflora* (lvs. next to stone) 200 m. on down on left, some large *Cibotium* and a clump of *Pandanus* *atleyanus* and *Clidemia*. Just beyond, on left, *Euphorbia* cf. *rockii* in tangle of *Gleichenia*, etc. Steep slope below with *Gleichenia* with *Gouldia* *term. v. crissata*, *Pilea* cf. *parviflora*, *Bobea elatior*, and down 300 m. *Peperomia mollis*. Beyond this a small landslide near with *Nephrolepis bicutata*. *Andropogon* along trail, then down to right *Phyllostegia grandiflora* & *Broussaisia* *Bobea elatior* on right, behind it a lehua with *Lycopodium phyllanthum* on lower trunk. 150 m down a large glaucous *Pouteria*. Then on left *Gleichenia glauca*. Down below on ridge top *Psychotria* cf. *haduana*.

large tree, below on right an area of typical tree forest.

Just beyond, some *Gleichenia* and some *Elaeocharis*.

Low part of trail - much *Psidium* etc. *Clidemia*, etc.

Several hundred yards along just before another *Gleichenia* slope -

Pittosporum sp.

on left and *Tetraploca* on right *Pittosporum* *gambelii*.

Lycopodium complanatum
Wolstenholmeia falconeri?
Leucophaea mollis

Under a large mossy leaning tree and around a curve a *Robea* tree

with *Leptocarpus* plate along right over trail on it.

Trees have epiphytic *Grammitis* *Elaph. rel.*

Well beyond this large curve another *Gleichenia* slope and excellent view of gulch.

Epiphytes abundant *Polypodium tamariscinum* several *Elaph.*

Great slopes of *Gleichenia* with large *Robea* trees and *Rob.*

Just before enormous tree a bush on right of *Clematis* *peruviana*.

~~*Hymenophyllum*~~ *Taraxacum* *boerhaavii* *Dorstenia* on left under *Cibotium* etc.

Deep ravine or cut through ridge to left.

Thunbergia bare cliff with *Cladonia* *Sphaerocarpon*, etc.

100 yds. beyond on rt. a hybrid *Leucophaea mollis* x *gambelii* like found. purple fls.

At a sharp left turn, on right *Myrsine* ~~with~~ *Leucophaea* in small *Gleichenia* patch scattered along trail.

Eupatorium *rupicola*.

On left at turn *Tournefortia* on left, *Pipturus* on rt.

Blechnum occidentale on bank on left.

Native bananas in gulch below.

Well along, on left a small ~~open~~ recess with a trunk with spectacular display of *Elaphoglossum* sp. + *Polypodium spectrum*.

100 m. on mt. *Heperomannia*
and a tree with *Grammitis*
and *Polykodium tamense*
Psilotum sp.

Then go on, a *Dubautia*
Plantago over head
Gouldia term. s. term. on mt.
and fairly common,
Well on up trail *Paleas*
become common. *Platyderma*
appears.

Acacia has remains a
prominent component
even more common locally
than *Metrosideros*

Broussaisia fairly common
Forest generally at this
alt. rather degraded.
open and filled with *Gleichenia*
Hymenophyllaceae on trees.

A great new landslide seen
below trail. Beyond this
the forest is in better condition,
but still not very good.
Less *Acacia* here. *Broussaisia*
abundant.

Along trail *Parhaleum conjugatum*
Centella, *Psidium* on cattl.

Forest of *Eugenia*, *Metrosideros*
Agrostoides, *Acacia*, *Gouldia*,
Perrottetia

At hairpin curve *Leptospermum*
scoparium (?) on ridge.

Little way along on
mt. below is a *Polypodium*
(only pinnate lfd. ~~tree~~)

Palea abundant

Chenodendron var.

Cibotium *eugenioides* &

C. splendens (?) side by side
on right.

Coprosma longifolia
hanging over trail in
left corner.

Some large *Chenodendron*
on mt.

Across valley of mt.

Clumps of *Brickellia*
Large new landslide seen

Then crossover to other side
of ridge. *Mechiaeria*
angustifolia common.

Midens sp. on cliff.

Leptospermum abundant
planted and ferns,
tree stand locally

Trachne distachyloides
on banks. *Cibotium*

glances on left.

Dubautia laxa overhead.

Trail goes up ridge.

Veg. dwarfed and scrubby.

Eugenia, *Psychotria*, *Palea* 2 spp.
Gouldia, *Myrsine* 2 spp.

Metrosideros, *Antidesma*
but *Leptospermum* is
very abundant.
Rye-cernuum, *Gleichenia*,

Pass over to rt. side of ridge
then on top again. *Phyllostegia*
grandiflora, *Dubautia*, *Bidens*.
Then to left side.

Large tree with 2 sp. *Hymen-*
phyllum, *Polypod. torn.*, *Etaph-*
etic. abundant mosses &
hepatia, *Hym. obtusum*, & *recurvata*
Wikstroemia, *Gouldia*,
Metrosideros, *Cheimodendron*
fabroea, *Machaerina ang.*
Dubautia laxa, *Cyanea*,
Sadleria cyathoides, *Gleichen-*
glauca, *Leptocarpus mollis*,
Coprosma longifolia,
Phyllostegia grand., *Cibotium*
glaberrimum, *Myrsine*
Psychotria, *Antidesma*
Broussaisia, *Freyerchia*,
Glech. linearis
all make up dense 2-3 m
scrub on high slopes.

Eub. inf. clear up here.

Tetrapharandra on left
A few *Ternstroemia* about
trail

This now seems fairly
good cloud forest.

More *Leptospermum*
Ternstroemia
Pritchardia next to
trail on left.
Cheimodendron, *Wikstroemia*
Pritchardia over trail.
Schizaea on vertical bank
Phyllostegia lanternaoides also.

Right turn into a ravine
with cloud scrub -
Dubautia, *Cibotium*, *Rabordia*,
Metrosideros, *Gouldia*,
Coprosma, *Broussaisia*,
Sadleria, *Bidens*, *Eugenia*,
Pelea, *Myrsine*.

Birouac spot.

Summit trail on other
side of ravine.

Turn left. 210 m.
Grassy marsh in
flat bottom of draw.

Scrub very depressed.
Lycopodium auratum
in boggy spots on left.
Sadleria hillebrandii
much *Machaerina*.

Gap at head of grassy
draw is badly trampled
and converted to grass.

Centella, Erechtites, etc.
On crest *Vaccinium dentatum*
near road, *Borreria*
Wikstroemia, *Gouldia*
Parbentia, *Cheirandrium*,
Eleocharis & *Peleocharis* sp.
Tetraplasandra, *Platydesma*,
Fabodia.

Low dense scrub
Down over edge, *Pithecharidia*
Cyperus, *Parbentia*,
Maclurea, *Bidens*,
Mitrosideros (rugosa?), *Gouldia*,
Phaneromeris, *Phyllostegia*
grand. - cont. *Trematolobelia*,
Blagayella menziesii
Gallieria cyathoides &
G. hillebrandii - *Pipturus*
Eupatorium riparium
Hedyotis centranthoides
Labordia, *Gouldia stipitata*,
Zanthoxylum.

Panicum, *Meritensia*, *Plantago*,
Racopilum, *Bidens*, *Gallieria*
hilleb. on boggy edge.

March 9. Head of Puunahua Valley
just north of head of Poamoho Trail
on wet windswept crest with
low scrubby vegetation.

- 41443 *Vittharia robusta*
5 rare, in shelter of bushes,
terrestrial
- 2 44 *Eugenia ~~sp.~~ koolauensis* Deg.
occasional
- 2 45 *Labordia*
rare
- 2 46 *Cyanea*
rare
- 3 47 *Coprosma longifolia*
common
- 1 48 same
- 1 49 *Broussaisia arguta* f. *arguta*
shrub 1 m. tall
- 3 50 *Cupatorium reparium*
occasional along trail
- 2 51 *Pipturus albidus* (H. & A.) Gray
common
- 4 52 *Vaccinium dentatum* Sm.
occasional

March 9. - Poamoho Trail
near Koolau crest

- 1 53 *Schizaea robusta*
occasional on vertical mossy bank along trail
- 2 54 *Lindsaea sp.*
in meadow-like forest
- 2 55 *Cladonia*
abundant on cut banks

800 m.

erect, caespitose

shrub 0.6 m. tall, fruit red.

shrub 1 m. tall, flowers yellow.

fleshy stemmed shrub 0.7 m.
tall, fruit yellow, fleshy.shrub 1 m. tall, fruit
immature, fls. pale green.
same 1.5 m. tall.flowers pale green with
rose purple stigmas.

weak shrub; flowers white.

shrub 1.5 m. tall; leaves
very rugose above, veins
red beneath, anthers green
& white.shrub 0.5 m. high, much
branched; flowers red with
green tips, fruit red.

775 m.

600 m.

March 10 - Manoa Cliff Trail
(from Round Top Dr. end)

Row of *Cordyline terminalis*
on left of entrance.

On trail proper, a strangling
fig (*Ficus aurea*?) on an
Acacia koa overhead.

Some large *Acacia koa* overhead
and *Aleurites moluccana*
on right. Trail grown up with
Setaria palmifolia.

about 100 m. from entrance
Freyrinetia on rt., *Persea*
overhead. 2 introduced *Elaeagnus*
above on left. *Ageratum bougainville*
in trail. *Emilia javanica*, *Puellia*
gracilis, *Passiflora edulis*,
Stachytarpheta sp.

Koa and *Kuhni* overhead.
About 100 m. more an extensive
bamboo grove. beyond this
a *Kuhni* forest on rt. *Pantanus*
on trail and *Rubus rosaeifolius*.
Bananas on right.

At first bend - some small
Cibotium ^{above} on left.

Commelina diffusa on small
flat place. Then *Guava* on
below. Strawberry *Guava* on left.
2 *Cibotium* on rt. *Jussiaea* in
trail. Strawberry *Guava*
becomes abundant 100 m
beyond bend. *Gleichenia*
lin. on left just before second
bend.

take left fork at this bend.
thru strawberry *Guava* brush.
Stachytarpheta on rt. here is
Grassy slopes - *Paspalum* sp.
Sacciolepis, *Setaria geniculata*,
Cassia bush. *Stachytarpheta*
bush for considerable dist
with *Koa*, *Freyrinetia*.

This is modified *Koa* forest.
Coming out of strawberry
Guava one encounters a
patch of *Cordyline*,
with several large *Guava*
above trail and *Kuhni* on rt.

From here trail lined
with *Cupatorium*. Some
Guava brush with *Cordyline*.
Then at a sharp left turn, first
small *Lehua*, *Lehua*
a scrub below trail, also *Cup.*
multiformis and *Lehua*,
native *Bidens*.

Open mixed scrubby wood
of *Lehua* & *Koa*. Just beyond
a large *Koa* hanging over
road *Pilea elatior* on right
and a thicket of *Psych.* (*strawberry*)
of *Kaduna*. Around a small
bend, under an *Aleurites* tree
several *Cyanea inquisititia*
mostly below trail. A little
farther, a slender *Coprosma long.*
Eragrostis variabilis
begins to appear on bank to left.

50 m. beyond bend *Scaevola gaudichaudiana* above trail among slender *Metrosideros* - these showing variation, narrow lfd. forms here. Just beyond this a large rock outcrop, then *Hebe* *arnottiana* below trail several trees. Beyond this clump, several bushes of *Coprosma foliosa* - neat, compact, narrow leaved. *Helaginella* many, common on rocks above trail, *Hebe* *communis*. Occ. *Spathoglottis*.

A little way on an open lava stand with *Freyinetia* ground cover of *Seteria* palm. *Guava* scrub below trail. Several *Bobes* slatins below trail. This *koa* stand continues for some distance with scattered *lehua*, *Scaevola*, *Psychotria*, *Guava*, strawberry *Guava* etc. Frosty trees below are *Kuhui*. Sharp bend to left - *Cragwort* *variabilis* on rocks. *Bidens*, also *Sphenomerus*. *Blechnum*, *Psidium* large *lehua* with curled lvs. below trail. Mixed scrub on steep cliff-like slope. Some *Lycopodium* *cernuum* in open ~~at~~ grassy spots. Around ~~at~~ bend is

at bushes of *Hylomeia* *hillebr.* on right, then a well-formed tree of *Eugenia* *sandwicensis*. Thicker *koa* forest with *Guava*, ^{strawberry} *Guava*, *Lantana*, *Freynetia* brush beneath. *Psychotria* up the slope, *Scaevola* common. Occasional white fld. *Lantana* below.

Then, on bend to left strawberry *Guava* shrubs with *koa* trees. *Lehua* below. *Scaevola* above. Beyond this *Pasiflora* *edulis* vines in opening of *Seteria* palm. View of *Kuhui* forest below. Along curve to rt & left open *koa* forest with large trees much *Freynetia*, *Palms* *rosaceifolius* & *Seteria* form ground cover.

After a couple of hundred m. to right, the trail bends sharply left. Just before this a *lehua* tree leans over trail with a clump of epiphytic *Nephrolepis* and wisps of *Vittaria*. If seen here mostly *koa*, open with *lehua*, some *Guava* and patch of *Kuhui* below. For 100 m then *bambos* *Guava* above trail. Then a patch of *lehua*, with *Scaevola* and *Psychotria*

Then *koa*. again,
with, below trail, several
Pouteria (coppery young growth)
also a small *Antidesma* plant.

A little way on an
ent. *melastom* & *Perrotetia*
on left, *Athyrium proliferum*?

Then large *quava* trees &
Cordyline. Beyond *quava*
Antidesma tree with
Dioscorea bulbifera & small
bananas below trail. A few
yards on *Glaucocarpus bifidus*
below trail. Then curve to left
with patch of *lehua*, *koa* &
Glaucocarpus. *Perrotetia*
hanging over trail, then
quava forest.

Through this, a curve from
which upper *manoa* can be
viewed, and *Konahuanui* if clear,
is *Psych. maritima* scrub
on right. A bit beyond this a
small *Gouldia* term. var. *curiosa*
on it at edge of cliff. Then
Diospyros ferrea v. *sanderi* also
on cliff. Open cliff with
Cordyline. Mixed scrub above
with *Dubautia plantaginea*.

In a small clump of wood
is *Hibiscus*, *Perrotetia*, *koa*,
Pipturus albidus. Then more
Cordyline on cliff. *Zingiber*
zerumbet in trail.

Next small clump of
wood is *Hibiscus* & *lehua*,
with *Paederia foetida*.

Then *quava* thicket
with some native plants.
Gouldia, *Hedyotis acuminata*,
Perrotetia, *Hibiscus*. In first
small opening, *Charpentiera*
above trail a few m.

Beyond next small
opening, *Osmanthus* sandw.
tree on it. mixed with
Citharexylum which looks
almost same sterile.

Sharp turn to left, then
mixed *koa*, *lehua*, *Hibiscus*,
with much *quava*. Around
a curve near *Cyanea*
angustifolia above trail.

A bit farther, at a small
bridge, in a *quava* thicket
some small *Wisonia* sh.

Then *koa* with *Euphorbia*,
Hibiscus, *Perrotetia*, *Gouldia*,
Cordyline, *Psychotria*, *Persea*,
etc. beneath.

Open slope with *Cordyline*
lehua & other bushes, much
Eupatorium. At other end
a little *Eupatorium adenophorum*.

Then *quava* thicket, some
native plants, much *Cordyline*.
Considerable walk thru
this, then several small

1961 Hawaiian Is.

Banana plants and
 Touchardia, on left. Fern
 covered rocks above with
 Adiantum, Selaginella
 + large hepatic. Embelia
 hanging over trail.

Wide ravine with
 Touchardia, bananas,
 cordylines, Coix, Colocasia,
 and Cyrtandra cf. cordifolia
 Prob. old Hawaiian cultivation.

Guava forest. Then
~~small~~ opening with
 Freziera, Ipomoea
 alba and Clermontia
 bahiana, this at base
 of large leaning koa on mt.
 Plex anomala just above
 on left. Several more
 Clermontia on mt.

Much Freziera in open
 koa wood. Much Hibiscus.
 Large Pisonia on small wooded
 ridge top to left.

From here on more lehua
 than koa.

Open slope with views of
 Mauna + mts. brush patches
 conspicuous. Arboretum
 planted forest in valley below.
 Brushy cliffs. Lehua +
 hibiscus with both Eupatoriums

Patch of wood with
 koa, Hibiscus, Antidesma,

lehua, cordylines, Citharexylum,
 around sharp curve + down
 (lehua). Dead koa on mt
 well epiphytic Nephrolepis
 exaltata. at next small
 curve Psychotria hexandra
 bush to it. Large membranous stipules

Guava-hibiscus forest.
 Athyrium proliferum patch
 on left.

Guava-lehua forest.
 Down slope into ~~the~~ lehua and
 lehua-koa forest, with
 Plex anomala, Clermontia,
 Perrottetia + Fouldia. Citharexylum
 abundant. Antidesma +
 Psychotria (Straussia)

Main trail. (Pt. to Pauoa
 down slope through koa-guava)
 Take left fork, then immediately
 another fork - left goes up Tantalus,
 straight ahead around Tantalus.
 Take this. Through ~~lehua~~ forest
 with ~~lehua~~ guava. Panakani
 esp. adrophorum. some lehua

Changes very soon to almost
 pure guava. with ground
 cover of Commelina diffusa.

Occasional patches of lawn,
 and Coix. Switchbacks.

Blechnum on banks.
 Nephrolepis cf. cordifolia or exaltata

abundant

Boulders on outcrops on left have
Trichomanes saxifragoides, *Adiantum* var.

Patches of *Pedicularis* (*coronarium*
+ *flavum*). *Thelypteris* *parvifolia*

Rejoin Pauoa trail.

Polypodium thurbergiae + *Peperomia*
obliqua on large rock, also *Trich. sax.*

Adiantum on mossy rocks on left
sudden change - lava to black sand.

Helopeltis abundant in black sand.

Kuhui forest in ^{open} *ho*.

with *lehua*, *Glacostegia*, *Flax*

Psychotria, *Antidesma*, *Samolites*

Freyeria, etc. on one straight
stretch of trail. then more

quava forest. *Persea*, etc. planted

in it. large kuhui +

various planted trees. then

thickets of coffee.

Stretch of quava forest, then

kuhui forest with planted *Albiggia*

falcata (?), then kuhui in draws,

quava on ridges, *oer. ho*, *lehua*,

kuhui in quava forest below to it.

Some areas have *Pipturus albidus*

Pagerstroemia planted, scattered

ho.

Eugenia jambos

Views of airport + Pearl Harbor, Barber Pt.

over Pauoa V. Honolulu Harbor

many exotic trees planted along
last part, incl. *Persea*, *Croton*, *Ficus*

Acacia, etc. *Heliconia* till. on road.

1 1/2 mi. bet.

entrance,

much *Hib. til.* on upper
Round Top Dr.

+ great display of exotics.

Round top slopes covered

by *Leucaena* scrub.

Black sand quarry

on it. on way up, fine

bedded black lapillaceous

tuff or "black sand"

